

Contents

| | | |
|----------|-------------------------------------------------------------|-----------|
| 1 | Introduction to Meiobenthology | 1 |
| 1.1 | Meiobenthos and Meiofauna: Definitions | 1 |
| 1.2 | A History of Meiobenthology | 2 |
| 2 | The Biotope: Factors and Study Methods | 7 |
| 2.1 | Abiotic Factors (Sediment Physiography) | 7 |
| 2.1.1 | Sediment Pores and Particles | 7 |
| 2.1.2 | Granulometric Characteristics | 9 |
| 2.1.3 | The Sediment–Water Regime | 14 |
| 2.1.4 | Physicochemical Characteristics | 22 |
| 2.2 | Biotic Habitat Factors: A Connected Complex | 37 |
| 2.2.1 | Detritus and Particulate Organic Matter (POM) | 38 |
| 2.2.2 | Dissolved Organic Matter (DOM) | 40 |
| 2.2.3 | Mucus, Exopolymers, and Biofilms | 41 |
| 2.2.4 | Bacteria | 43 |
| 2.2.5 | Microphytobenthos | 48 |
| 2.2.6 | Higher Plants | 53 |
| 2.2.7 | Animals Structuring the Ecosystem | 53 |
| 2.3 | Conclusion: The Microtexture of Natural Sediments | 59 |
| 3 | Sampling and Processing Meiofauna | 63 |
| 3.1 | Sampling | 63 |
| 3.1.1 | Number of Replicates and Size of Sampling Units | 63 |
| 3.1.2 | Sampling Devices | 64 |
| 3.2 | Processing of Meiofaunal Samples | 72 |
| 3.2.1 | Preserving Meiofauna in Their Natural Void System | 72 |
| 3.2.2 | Extraction of Meiofauna | 73 |
| 3.2.3 | Fixation and Preservation | 77 |
| 3.2.4 | Processing and Identifying Meiofaunal Organisms | 80 |
| 3.3 | Extraction of Pore Water | 84 |

| | | |
|----------|---------------------------------------------------------------------------------------------------------------------|-----|
| 4 | Biological Characteristics of Meiofauna | 87 |
| 4.1 | Adaptations to the Biotope | 87 |
| 4.1.1 | Adaptations to Narrow Spaces: Miniaturization, Elongation, Flexibility | 87 |
| 4.1.2 | Adaptations to the Mobile Environment: Adhesion, Special Locomotion, Reinforcing Structures | 92 |
| 4.1.3 | Adaptations to the Three-Dimensional Dark Environment: Static Organs, Reduction of Pigment and Eyes | 97 |
| 4.1.4 | Adaptations Related to Reproduction and Development | 99 |
| 5 | Meiofauna Taxa: A Systematic Account | 103 |
| 5.1 | Protista (Protoctista) | 103 |
| 5.1.1 | Foraminifera (Rhizaria: Granuloreticulosa) | 103 |
| 5.1.2 | Heliozoa (Actinopodia). | 107 |
| 5.1.3 | Amoebozoa (“Rhizopoda”): Gymnamoebia, Testacea. | 107 |
| 5.1.4 | Ciliophora (Ciliata). | 108 |
| 5.2 | Cnidaria. | 114 |
| 5.2.1 | Hydrozoa (Medusae). | 116 |
| 5.2.2 | Hydrozoa (Polyps). | 116 |
| 5.2.3 | Scyphozoa. | 118 |
| 5.2.4 | Anthozoa. | 118 |
| 5.3 | Free-Living Platyhelminthes: Turbellarians | 119 |
| 5.3.1 | Major Turbellarian Groups | 120 |
| 5.3.2 | Distributional and Ecological Aspects | 123 |
| 5.4 | Gnathifera | 127 |
| 5.4.1 | Gnathostomulida. | 127 |
| 5.4.2 | Rotifera, Rotatoria | 129 |
| 5.4.3 | Micrognathozoa | 133 |
| 5.5 | Nemertinea | 134 |
| 5.6 | Nemathelminthes: A Valid Taxon? | 136 |
| 5.6.1 | Nematoda (Free-Living). | 137 |
| 5.6.2 | Kinorhyncha | 156 |
| 5.6.3 | Priapulida | 158 |
| 5.6.4 | Loricifera | 160 |
| 5.6.5 | Gastrotricha. | 162 |
| 5.7 | Tardigrada | 165 |
| 5.8 | Crustacea. | 171 |
| 5.8.1 | Cephalocarida | 172 |
| 5.8.2 | Anostraca: Anomopoda (“Cladocera”; “Branchiopoda”). | 173 |
| 5.8.3 | Ostracoda | 175 |

| | | |
|----------|-----------------------------------------------------------------------------------|------------|
| 5.8.4 | Mystacocarida | 180 |
| 5.8.5 | Copepoda: Harpacticoida | 181 |
| 5.8.6 | Copepoda: Cyclopoida and Siphonostomatoida | 189 |
| 5.8.7 | Malacostraca | 190 |
| 5.9 | Chelicerata: Acari | 201 |
| 5.9.1 | Halacaroida: Halacaridae | 201 |
| 5.9.2 | Freshwater Mites: "Hydrachnidia," Stygothrombiidae, and Others | 205 |
| 5.9.3 | Palpigradi (Arachnida) | 205 |
| 5.9.4 | Pycnogonida, Pantopoda | 206 |
| 5.10 | Terrigenous Arthropoda (Thalassobionts) | 207 |
| 5.11 | Annelida | 207 |
| 5.11.1 | Polychaeta | 208 |
| 5.11.2 | Oligochaeta | 215 |
| 5.11.3 | Annelida "Incertae sedis" | 218 |
| 5.12 | Sipuncula | 221 |
| 5.13 | Mollusca | 223 |
| 5.13.1 | Monoplacophora and Aplacophora | 223 |
| 5.13.2 | Gastropoda | 225 |
| 5.14 | Tentaculata | 226 |
| 5.14.1 | Brachiopoda | 226 |
| 5.14.2 | Bryozoa, Ectoprocta | 227 |
| 5.15 | Kamptozoa, Entoprocta | 228 |
| 5.16 | Echinodermata | 229 |
| 5.16.1 | Holothuroidea | 229 |
| 5.17 | Chaetognatha | 230 |
| 5.18 | Tunicata (Chordata) | 231 |
| 5.18.1 | Asciacea | 231 |
| 5.18.2 | Sorberacea | 232 |
| 5.19 | Meiofaunal Taxa: Concluding Remarks | 233 |
| 6 | Evolutionary and Phylogenetic Effects in Meiobenthology | 235 |
| 6.1 | Body Structures of Evolutionary Relevance | 235 |
| 6.2 | Meiofauna in the Fossil Record | 239 |
| 7 | Patterns of Meiofauna Distribution | 243 |
| 7.1 | Evolutionary Aspects | 243 |
| 7.2 | Zoogeographic Aspects | 249 |
| 7.2.1 | Mechanisms of Dispersal | 250 |
| 7.2.2 | Geological Structures and Processes | 256 |
| 7.3 | Ecological Aspects of Distributional Importance: Horizontal Patterns | 259 |
| 7.4 | Vertical Zonation of Meiobenthos | 261 |

| | | |
|----------|-------------------------------------------------------------------------------------------------|-----|
| 8 | Meiofauna from Selected Biotopes and Regions | 267 |
| 8.1 | Polar Regions | 268 |
| 8.1.1 | Sea Ice | 270 |
| 8.2 | Marine Subtropical and Tropical Regions | 276 |
| 8.2.1 | Tropical Sands | 278 |
| 8.2.2 | Mangroves | 280 |
| 8.3 | The Deep-Sea | 284 |
| 8.3.1 | The Habitat | 284 |
| 8.3.2 | The Meiofauna | 287 |
| 8.4 | Dysoxic, Anoxic, and Sulfidic Environments: Discussing the Thiobios | 296 |
| 8.4.1 | Reducing Habitats of the Thiobios | 296 |
| 8.4.2 | Thiobiotic Meiobenthos | 298 |
| 8.4.3 | Survival of Thiobios Under Anoxia and Sulphide – Mechanisms and Adaptations | 302 |
| 8.4.4 | Food Spectrum of the Thiobios | 307 |
| 8.4.5 | Distribution and Succession of the Thiobios | 308 |
| 8.4.6 | Diversity and Evolution of the Thiobios | 309 |
| 8.4.7 | Chemoautotrophy-Based Ecosystems: Vents, Seeps, and Other Exotic Habitats | 313 |
| 8.5 | Phytoplankton Habitats and Hard Substrates | 317 |
| 8.6 | Brackish Water Sites | 324 |
| 8.7 | Freshwater Biotopes | 328 |
| 8.7.1 | Running Waters: Stream and River Beds | 329 |
| 8.7.2 | The Groundwater System | 338 |
| 8.7.3 | Standing Waters, Lakes | 344 |
| 8.8 | Polluted Habitats | 349 |
| 8.8.1 | General Aspects and Method Survey | 349 |
| 8.8.2 | Selected Cases of Pollution and Meiofauna | 361 |
| 9 | Synecological Perspectives in Meiobenthology | 373 |
| 9.1 | Community Structure and Diversity | 373 |
| 9.1.1 | Processes of Recolonization | 375 |
| 9.2 | Community Structure and Size Spectra | 377 |
| 9.3 | The Meiobenthos in the Benthic Energy Flow | 383 |
| 9.3.1 | General Considerations | 383 |
| 9.3.2 | Assessing Production: Abundance, Biomass, P/B Ratio, Respiration | 387 |
| 9.3.3 | The Energetic Divergence Between Meiofauna and Macrofauna | 397 |
| 9.4 | The Position of Meiofauna in the Benthic Ecosystem: A Compilation of Energy Fluxes | 400 |
| 9.4.1 | The Meiofauna as Members of the “Small Food Web” | 402 |
| 9.4.2 | Links Between the Meiofauna and the Macrofauna | 406 |
| 9.4.3 | Meiofauna as an Integrative Benthic Complex | 410 |

| | |
|--------------------------------------------------------------------------------------------------------|------|
| Contents | xvii |
| 10 Retrospect on Meiobenthology and Outlook on New Approaches and Future Research | 417 |
| References | 423 |
| Glossary | 503 |
| Index | 513 |



<http://www.springer.com/978-3-540-68657-6>

Meiobenthology

The Microscopic Motile Fauna of Aquatic Sediments

Giere, O.

2009, XVII, 527 p., Hardcover

ISBN: 978-3-540-68657-6