

Americans undergo colonoscopies too often, study finds

Following the proper guidelines could save money and discomfort of unnecessary screenings

Colonoscopies are a very valuable procedure by which to screen for the presence of colorectal cancer. However, it seems that healthy Americans who do undergo this sometimes uncomfortable examination often have repeat screenings long before they actually should. Gina Kruse of Massachusetts General Hospital in the US and colleagues advise that endoscopists stick to the national guidelines more closely. Their findings¹ appear in the *Journal of General Internal Medicine*², published by Springer.

Current national guidelines strongly recommend that adults aged 50 and older should be screened every ten years, while surveillance colonoscopies should be performed more frequently in adults with pre-cancerous polyps called adenomas. Concern has been raised that these guidelines are not being followed. Kruse's research team therefore set out to measure if screening and surveillance colonoscopies among average-risk adults are really overused, and why this might be so. They used electronic health record data of 1,429 adults between the ages of 50 to 65 years old who underwent an initial colonoscopy for cancer screening between 2001 and 2010. Subsequent exams were done in 871 cases, on average six years after the initial procedure.

They found that 88 percent of the follow-up screening colonoscopies and one in every two surveillance colonoscopies (49 percent) were done earlier than guidelines recommend. People who had no signs of abnormal growths such as polyps the first time around were again examined on average 6.9 years later. Cases where benign polyps (non-adenomas) were found were re-examined on average 5.9 years later.

The researchers say this means that patients who are enrolled in a screening program from the age of 50 to 74 years actually undergo one additional colonoscopy in their lifetimes compared to what is recommended by current guidelines. On the flipside, one in every four exams in which pre-cancerous tumors were initially detected was not followed up within the recommended guideline interval.

The researchers found a strong association between endoscopist recommendations and early follow-up interval for colonoscopy, consistent with prior research indicating many endoscopists do not agree with the follow-up intervals recommended in the guidelines. They therefore advise that healthcare systems consider monitoring follow-up recommendations as an important lever for discussion and potential reduction in overuse of colonoscopy.

Senior study author Thomas Sequist of Brigham and Women's Hospital in the US said, "Examining practice variation and establishing locally endorsed standards among endoscopists may be a way to target interventions to reduce overuse."

"The idea that a large proportion of the 14 million screening colonoscopies performed annually in the US are actually done unnecessarily is especially concerning in light of the 28 million Americans between 50 and 74 who are not up to date in colorectal cancer screening," says Gina Kruse. "The overused colonoscopies on the patients in this study alone represent a potential excess of over \$1 million in health care spending—resources that might benefit those who are overdue for colon cancer screening."

References:

1. Kruse, G.R. *et al* (2014). Overuse of Colonoscopy for Colorectal Cancer Screening and Surveillance, *Journal of General Internal Medicine*. DOI 10.1007/s11606-014-3015-6
2. The *Journal of General Internal Medicine* is the official journal of the Society of General Internal Medicine.

The full-text article is available to journalists on request.

Contact: Joan Robinson | Springer | tel +49-6221-487-8130 | joan.robinson@springer.com



<http://www.springer.com/journal/11606>

Journal of General Internal Medicine
Editor-in-Chief: Kravitz, R.; Feldman, M.
ISSN: 0884-8734 (print version)
ISSN: 1525-1497 (electronic version)
Journal no. 11606