
Cyclic Homology with Coefficients

D. Kaledin*

Steklov Math Institute, Moscow, Russia
kaledin@mccme.ru

To Yu. I. Manin, the founder, on the occasion of his 70th birthday

Summary. We propose a category which can serve as the category of coefficients for the cyclic homology $HC_*(A)$ of an associative algebra A over a field k . The construction is categorical in nature, and essentially uses only the tensor category $A\text{-bimod}$ of A -bimodules; objects of our category are A -bimodules with an additional structure. We also generalize the construction to a more general tensor k -linear tensor category C instead of $A\text{-bimod}$. We add some remarks about Getzler's version of the Gauss–Manin connection for the periodic cyclic homology $HP_*(A)$.

Key words: cyclic homology, Gauss–Manin connection, tensor categories

2000 Mathematics Subject Classifications: 19D55, 18E25, 18G10

Introduction

Ever since it was discovered in 1982 by A. Connes [C1] and B. Tsygan [Ts], cyclic homology has occupied a strange place in the realm of homological algebra. Normally in homological algebra problems, one expects to start from some data, for instance a topological space X , then construct some abelian category, such as the category of sheaves on X , and then define the cohomology of X by computing the derived functors of some natural functor, such as the global sections functor $\Gamma(X, -)$. Admittedly, this is a modern formulation, but it was certainly current already in 1982. Cyclic homology starts with an associative algebra A , and defines its homology groups $HC_*(A)$, but there are absolutely no derived functors in sight. Originally, groups $HC_*(A)$ were defined as the homology of an explicit complex, which anyone trained to use triangulated categories cannot help but take as an insult. Later, A.

*Partially supported by CRDF grant RUM1-2694-MO05.

Connes [C2] improved on the definition by introducing the abelian category of so-called cyclic vector spaces. However, the passage from A to its associated cyclic vector space $A_{\#}$ is still done by an explicit *ad hoc* formula. It is as if we were to know the bar-complex that computes the homology of a group, without knowing the definition of the homology of a group.

This situation undoubtedly irked many people over the years, but to the best of my knowledge, no satisfactory solution has been proposed, and it may not exist; indeed, many relations to the de Rham homology notwithstanding, it is not clear whether cyclic homology properly forms a part of homological algebra at all (to the point that for instance in [FT], the word “homology” is not used at all for $HC_{\bullet}(A)$, and it is called instead *additive K-theory* of A). In the great codification of homological algebra done in [GM1], cyclic homology appears only in the exercises. This is not surprising, since the main unifying idea of [GM1] is the ideology of “linearization”: homological algebra linearizes geometry, just as functional analysis used to do 50 years ago; triangulated categories and adjoint functors are modern-day versions of Banach spaces and adjoint linear operators. This has been an immensely successful and clarifying point of view in general, but $HC_{\bullet}(A)$ sticks out on a complete tangent: there is simply no natural place for it in this framework.

This paper arose as one more attempt to propose a solution to the difficulty: to find a natural triangulated category where $HC_{\bullet}(-)$ would be able to live with a certain level of comfort (and with all the standard corollaries such as the notion of cyclic homology with coefficients, the ability to compute cyclic homology by whatever resolution is convenient, not just the bar resolution, and so on).

In a sense, our attempt has been successful: we define a triangulated category that can serve as the natural “category of coefficients” for cyclic homology of an algebra A , and we prove the comparison theorem that shows that when the coefficients are trivial, the new definition of cyclic homology is equivalent to the old one. In fact, the algebra A enters into the construction only through the category $A\text{-bimod}$ of A -bimodules; we also show how to generalize the construction so that $A\text{-bimod}$ is replaced with a more general tensor abelian category \mathcal{C} .

From a different point of view, though, our attempt failed miserably: the correspondence $A \mapsto A_{\#}$, being thrown out of the window, immediately returns through the door in a new and “higher-level” disguise: it is now applied not to the algebra A , but to the tensor category $\mathcal{C} = A\text{-bimod}$. Then in practice, the freedom to choose an arbitrary resolution to compute the derived functors leads, in our approach to $HC_{\bullet}(-)$, to complexes that are even larger than the original complex, and at some point the whole exercise starts to look pointless.

Still, we believe that all said and done, some point can be found, and some things are clarified in our approach; one such thing is, for instance, the version of Gauss–Manin connection for cyclic homology discovered by E. Getzler [Ge]. Moreover, we do propose a definition of cyclic homology

that – makes sense for a general tensor category; and in some particular questions, even the computations can be simplified. As for the presence of the $A_{\#}$ -construction, this might be in the nature of things, after all: not a bug in the theory, but a necessary feature. However, we leave it to the reader to be the judge.

The paper is organized as follows. In Section 1 we recall A. Connes’s second definition of cyclic homology, which uses the cyclic category \mathcal{A} ; we also recall some facts about homology of small categories that we will need. We have tried to give only the absolute minimum; the reader not familiar with the material will have to consult the references. In Section 2 we introduce our main object: the notion of a *cyclic bimodule* over an associative algebra A , and the derived category of such bimodules. We also introduce cyclic homology $HC_{\bullet}(A, M)$ with coefficients in a cyclic bimodule M . In Section 3 we give a very short derivation of the Gauss–Manin connection; strictly speaking, the language of cyclic bimodules is not needed for this, but we believe that it shows more clearly what is really going on. In Section 4, we show how to replace the category $A\text{-bimod}$ everywhere with a more general tensor abelian category \mathcal{C} . Section 5 is a postface, or a “discussion” (as is done in medical journals); we discuss some of the further things one might (and should) do with cyclic bimodules, and how to correct some deficiencies of the theory developed in Sections 2 and 4.

Acknowledgments.

In the course of this work, I have benefited greatly from discussions with A. Beilinson, E. Getzler, V. Ginzburg, A. Kuznetsov, N. Markarian, D. Tamarkin, and B. Tsygan. I am grateful to Northwestern University, where part of this work was done, and where some of the results were presented in seminars, with great indulgence from the audience toward the unfinished state of the work. And last but not least, it is a great pleasure and a great opportunity to dedicate this paper to Yuri Ivanovich Manin on his birthday. Besides all the usual things, I would like to stress that it is the book [GM1], and [GM2] to a lesser extent, that shaped the way we look at homological algebra today, at least “we” of my generation and of the Moscow school. Without Manin’s decisive influence, this paper certainly would not have appeared (as in fact at least half of the papers I have written).

1 Recollection on cyclic homology.

We start by recalling, extremely briefly, A. Connes’ approach to cyclic homology, which was originally introduced in [C2] (for detailed overviews, see, e.g., [L, Section 6] or [FT, Appendix]; a brief but complete exposition using the same language and notation as in this paper can be found in [Ka, Section 1]).

Connes' approach relies on the technique of homology of small categories. Fix a base field k . Recall that for every small category Γ , the category $\text{Fun}(\Gamma, k)$ of functors from Γ to the category $k\text{-Vect}$ of k -vector spaces is an abelian category with enough projectives and enough injectives, with derived category $\mathcal{D}(\Gamma, k)$. For any object $E \in \text{Fun}(\Gamma, k)$, the homology $H_*(\Gamma, E)$ of the category Γ with coefficients in E is by definition the derived functor of the direct limit functor

$$\lim_{\overline{\Gamma}} : \text{Fun}(\Gamma, k) \rightarrow k\text{-Vect}.$$

Analogously, the cohomology $H^*(\Gamma, E)$ is the derived functor of the inverse limit $\lim_{\overline{\Gamma}}$. Equivalently,

$$H^*(\Gamma, E) = \text{Ext}^*(k, E),$$

where $k \in \text{Fun}(\Gamma, k)$ is the constant functor (all objects in Γ go to k , all maps go to identity). In particular, $H^*(\Gamma, k)$ is an algebra. For any $E \in \text{Fun}(\Gamma, k)$, the cohomology $H^*(\Gamma, E)$ and the homology $H_*(\Gamma, E)$ are modules over $H^*(\Gamma, k)$.

We also note, although it is not needed for the definition of cyclic homology, that for any functor $\gamma : \Gamma' \rightarrow \Gamma$ between two small categories, we have the pullback functor $\gamma^* : \text{Fun}(\Gamma, k) \rightarrow \text{Fun}(\Gamma', k)$, and for any $E \in \text{Fun}(\Gamma, k)$, we have natural maps

$$H_*(\Gamma', \gamma^*E) \rightarrow H_*(\Gamma, E), \quad H^*(\Gamma, E) \rightarrow H^*(\Gamma', \gamma^*E). \quad (1.1)$$

Moreover, the pullback functor γ^* has a left adjoint $\gamma_! : \text{Fun}(\Gamma', k) \rightarrow \text{Fun}(\Gamma, k)$ and a right-adjoint $f_* : \text{Fun}(\Gamma', k) \rightarrow \text{Fun}(\Gamma, k)$, known as the left and right Kan extensions. In general, $f_!$ is right-exact, but it need not be left-exact. We will need one particular case in which it is exact. Assume given a covariant functor $V : \Gamma \rightarrow \text{Sets}$ from a small category Γ to the category of sets, and consider the category Γ' of pairs $\langle [a], v \rangle$ of an object $[a] \in \Gamma$ and an element $v \in V([a])$ (maps in Γ' are those maps $\gamma : [a] \rightarrow [a']$ that send $v \in V([a])$ to $v' \in V([a'])$). Such a category is known as a *discrete cofibration* over Γ associated to V ; see [Gr]. Then the Kan extension $f_!$ associated to the forgetful functor $f : \Gamma' \rightarrow \Gamma, \langle [a], v \rangle \mapsto [a]$ is exact, and is easy to compute: for any $E \in \text{Fun}(\Gamma', k)$ and $[a] \in \Gamma$, we have

$$f_!E([a]) = \bigoplus_{v \in V([a])} E(\langle [a], v \rangle). \quad (1.2)$$

Moreover, for any $E \in \text{Fun}(\Gamma, k)$, this immediately gives the projection formula:

$$f_!f^*E \cong E \otimes F_!k, \quad (1.3)$$

where as before, $k \in \text{Fun}(\Gamma', k)$ stands for the constant functor.

For applications to cyclic homology, one starts by introducing the *cyclic category* Λ . This is a small category whose objects $[n]$ are numbered by positive integers $n \geq 1$. One thinks of an object $[n]$ as a circle S^1 with n distinct marked points; we denote the set of these points by $V([n])$. The set of maps $\Lambda([n'], [n])$ from $[n']$ to $[n]$ is then the set of homotopy classes of continuous maps $f : S^1 \rightarrow S^1$ such that

- f has degree 1, sends marked points to marked points, and is nondecreasing with respect to the natural cyclic order on S^1 (that is, if a point $a \in S^1$ lies between points b and c when one counts clockwise, then the same is true for $f(a)$, $f(b)$, and $f(c)$).

In particular, we have $\Lambda([1], [n]) = V([n])$. This topological description of the cyclic category Λ is easy to visualize, but there are also alternative combinatorial descriptions (e.g., [GM1, Exercise II.1.6], [L, Section 6], or [FT, A.2], retold in [Ka, Section 1.4]). All the descriptions are equivalent. Objects in $\text{Fun}(\Lambda, k)$ are usually called *cyclic vector spaces*.

The cyclic category Λ is related to the more familiar *simplicial category* Δ^{opp} , the opposite to the category Δ of finite nonempty linearly ordered sets. To understand the relation, consider the discrete cofibration $\Lambda_{[1]}/\Lambda$ associated to the functor $V : \Lambda \rightarrow \text{Sets}$; equivalently, $\Lambda_{[1]}$ is the category of objects $[n]$ in Λ equipped with a map $[1] \rightarrow [n]$. Then it is easy to check that $\Lambda_{[1]}$ is equivalent to the category Δ^{opp} . From now on, we will abuse notation and identify $\Lambda_{[1]}$ and Δ^{opp} . We then have a natural projection $\Delta^{opp} = \Lambda_{[1]} \rightarrow \Lambda$, $\langle [n], v \rangle \mapsto [n]$, which we denote by $j : \Delta^{opp} \rightarrow \Lambda$.

For any cyclic k -vector space $E \in \text{Fun}(\Lambda, k)$, we have its restriction $j^*E \in \text{Fun}(\Delta^{opp}, E)$, a simplicial vector space. One defines the cyclic homology $HC_\bullet(E)$ and the Hochschild homology HH_\bullet of E by

$$HC_\bullet(E) \stackrel{\text{def}}{=} H_\bullet(\Lambda, E), \quad HH_\bullet(E) \stackrel{\text{def}}{=} H_\bullet(\Delta^{opp}, j^*E).$$

By (1.1), we have a natural map $HH_\bullet(E) \rightarrow HC_\bullet(E)$ (moreover, since $j : \Delta^{opp} \rightarrow \Lambda$ is a discrete cofibration, the Kan extension $j_!$ is exact, so that we have $HH_\bullet(E) \cong HC_\bullet(j_!j^*E)$, and the natural map is induced by the adjunction map $j_!j^*E \rightarrow E$). It has been shown by A. Connes that this map fits into a long exact sequence

$$HH_\bullet(E) \longrightarrow HC_\bullet(E) \xrightarrow{u} HC_{\bullet-2}(E) \longrightarrow \dots \quad (1.4)$$

Here the map u is the so-called *periodicity map* on $HC_\bullet(E)$: one shows that the algebra $H^*(\Lambda, k)$ is isomorphic to the polynomial algebra $k[u]$ in one generator u of degree 2, and the periodicity map on homology is simply the action of this generator. This allows one to define a third homological invariant, the *periodic cyclic homology* $HP_\bullet(E)$; to do it, one inverts the periodicity map.

Definition 1.1. For any cyclic k -vector space $E \in \text{Fun}(A, k)$, the periodic cyclic homology of E is defined by

$$HP_{\bullet}(E) = \varprojlim_{\bullet} HC_{\bullet}(E),$$

where \varprojlim_{\bullet} denotes the derived functor of the inverse limit \varprojlim .

Assume now that we are given an associative unital algebra A over k . To define its cyclic homology, we associate to A a canonical cyclic vector space $A_{\#}$ in the following way. We set $A_{\#}([n]) = A^{\otimes V([n])}$, the tensor product of n copies of the vector space A numbered by marked points $v \in V([n])$. Then for any map $f \in A([n'], [n])$, we define

$$A_{\#}(f) = \bigotimes_{v \in V([n])} m_{f^{-1}(v)} : A^{\otimes V([n'])} = \bigotimes_{v \in V([n])} A^{\otimes f^{-1}(v)} \rightarrow A^{\otimes V([n])}, \quad (1.5)$$

where for any linearly ordered finite set S , $m_S : A^{\otimes S} \rightarrow A$ is the canonical multiplication map induced by the associative algebra structure on A (and if S is empty, we set $A^{\otimes S} = k$, and m_S is the embedding of the unity). This is obviously compatible with compositions, and it is well defined since for any $v \in V([n])$, its preimage $f^{-1} \subset V([n'])$ carries a natural linear order induced by the orientation of the circle S^1 .

Definition 1.2. For any associative unital algebra A over k , its Hochschild, cyclic, and periodic cyclic homologies $HH_{\bullet}(A)$, $HC_{\bullet}(A)$, $HP_{\bullet}(A)$ are defined as the corresponding homologies of the cyclic k -vector space $A_{\#}$:

$$HH_{\bullet}(A) \stackrel{\text{def}}{=} HH_{\bullet}(A_{\#}), \quad HC_{\bullet}(A) \stackrel{\text{def}}{=} HC_{\bullet}(A_{\#}), \quad HP_{\bullet}(A) \stackrel{\text{def}}{=} HP_{\bullet}(A_{\#}).$$

2 Cyclic bimodules.

Among all the homology functors introduced in Definition 1.2, Hochschild homology is the most accessible, and this is because it has another definition: for any associative unital algebra A over k , we have

$$HH_{\bullet} = \text{Tor}_{A^{opp} \otimes A}^{\bullet}(A, A), \quad (2.1)$$

where Tor^{\bullet} is taken over the algebra $A^{opp} \otimes A$ (here A^{opp} denotes A with the multiplication taken in the opposite direction).

This has a version with coefficients: if M is a left module over $A^{opp} \otimes A$, in other words, an A -bimodule, one defines Hochschild homology of A with coefficients in M by

$$HH_{\bullet}(A, M) = \text{Tor}_{A^{opp} \otimes A}^{\bullet}(M, A). \quad (2.2)$$

The category $A\text{-bimod}$ of A -bimodules is a unital (nonsymmetric) tensor category, with tensor product $- \otimes_A -$ and the unit object A . Hochschild homology is a homological functor from $A\text{-bimod}$ to $k\text{-Vect}$.

To obtain a small category interpretation of $HH_*(A, M)$, one notes that for any $n, n' \geq 0$, the A -bimodule structure on M induces a multiplication map

$$A^{\otimes n} \otimes M \otimes A^{\otimes n'} \rightarrow M.$$

Therefore, if to any $\langle [n], v \rangle \in \Delta^{opp}$ we associate the k -vector space

$$M_{\#}^{\Delta}([n]) = M \otimes A^{\otimes (V([n]) \setminus \{v\})}, \quad (2.3)$$

with M filling the place corresponding to $v \in V([n])$, then (1.5) makes perfect sense for those maps $f : [n'] \rightarrow [n]$ that preserve the distinguished points. Thus to any $M \in A\text{-bimod}$, we can associate a simplicial k -vector space $M_{\#}^{\Delta} \in \text{Fun}(\Delta^{opp}, k)$. In the particular case $M = A$, we have $A_{\#}^{\Delta} = j^* A_{\#}$.

Lemma 2.1. *For any $M \in A\text{-bimod}$, we have a canonical isomorphism*

$$HH_*(A, M) \cong H_*(\Delta^{opp}, M_{\#}^{\Delta}). \quad (2.4)$$

Proof. It is well known that for any simplicial k -vector space E , the homology $H_*(\Delta^{opp}, E)$ can be computed by the standard complex of E (that is, the complex with terms $E([n])$ and the differential $d = \sum_i (-1)^i d_i$, where d_i are the face maps). In particular, $H_0(\Delta^{opp}, M_{\#}^{\Delta})$ is the cokernel of the map $d : A \otimes M \rightarrow M$ given by $d(a \otimes m) = am - ma$. The natural projection $M \rightarrow M \otimes_{A^{opp} \otimes A} A$ obviously factors through this cokernel, so that we have a natural map

$$\rho_0 : H_0(\Delta^{opp}, M_{\#}^{\Delta}) \rightarrow HH_0(A, M).$$

Both sides of (2.4) are homological functors in M , and $HH_*(A, M)$ is a universal homological functor (= the derived functor of $HH_0(A, M)$); therefore the map ρ_0 extends to a map $\rho_{\bullet} : H_*(\Delta^{opp}, M_{\#}^{\Delta}) \rightarrow HH_*(A, M)$. To prove that ρ_{\bullet} is an isomorphism for any M , it suffices to prove it when M is free over $A^{opp} \otimes A$, or in fact, when $M = A^{opp} \otimes A$. Then on the one hand, $HH_0(A, M) = A$, and $HH_i(A, M) = 0$ for $i \geq 1$. And on the other hand, the standard complex associated to the simplicial k -vector space $(A^{opp} \otimes A)_{\#}^{\Delta}$ is just the usual bar resolution of the diagonal A -bimodule A .

It is more or less obvious that for an arbitrary $M \in A\text{-bimod}$, $M_{\#}^{\Delta}$ does not extend to a cyclic vector space: in order to be able to define $HC_*(A, M)$, we have to equip the bimodule M with some additional structure. To do this, we want to use the tensor structure on $A\text{-bimod}$. The slogan is the following:

- To find a suitable category of coefficients for cyclic homology, we have to repeat the definition of the cyclic vector space $A_{\#} \in \text{Fun}(A, k)$, but replace the associative algebra A in this definition with the tensor category $A\text{-bimod}$.

Let us explain what this means.

First, consider an arbitrary associative unital monoidal category \mathcal{C} with unit object I (at this point, not necessarily abelian). For any integer n , we have the Cartesian product $\mathcal{C}^n = \mathcal{C} \times \mathcal{C} \times \cdots \times \mathcal{C}$. Moreover, the product on \mathcal{C} induces a product functor

$$m : \mathcal{C}^n \rightarrow \mathcal{C},$$

where if $n = 0$, we let $\mathcal{C}^n = \mathbf{pt}$, the category with one object and one morphism, and let $m : \mathbf{pt} \rightarrow \mathcal{C}$ be the embedding of the unit object. More generally, for any finite linearly ordered set S with n elements, we have a product functor $m_S : \mathcal{C}^S \rightarrow \mathcal{C}$, where $\mathcal{C}^S = \mathcal{C}^n$ with multiples in the product labeled by elements of S . Then for any $[n], [n'] \in \Lambda$, and any $f : [n'] \rightarrow [n]$, we can define a functor $f_! : \mathcal{C}^{V([n'])} \rightarrow \mathcal{C}^{V([n])}$ by the same formula as in (1.5):

$$f_! = \prod_{v \in V([n])} m_{f^{-1}(v)} : \mathcal{C}^{V([n'])} = \prod_{v \in V([n])} \mathcal{C}^{f^{-1}(v)} \rightarrow \mathcal{C}^{V([n])}. \quad (2.5)$$

The natural associativity isomorphism for the product on \mathcal{C} induces natural isomorphisms $(f \circ f')_! \cong f_! \circ f'_!$, and one checks easily that they satisfy natural compatibility conditions. All in all, setting $[n] \mapsto \mathcal{C}^{V([n])}$, $f \mapsto f_!$ defines a weak functor (a.k.a. Lax functor, a.k.a. 2-functor, a.k.a. pseudofunctor in the original terminology of Grothendieck) from Λ to the category of categories. Informally, we have a “cyclic category.”

To work with weak functors, it is convenient to follow Grothendieck’s approach in [Gr]. Namely, instead of considering a weak functor directly, we define a *category* $\mathcal{C}_\#$ in the following way: its objects are pairs $\langle [n], M_n \rangle$ of an object $[n]$ of Λ and an object $M_n \in \mathcal{C}^n$, and morphisms from $\langle [n'], M_{n'} \rangle$ to $\langle [n], M_n \rangle$ are pairs $\langle f, \iota_f \rangle$ of a map $f : [n'] \rightarrow [n]$ and a bimodule map $\iota_f : f_!(M_{n'}) \rightarrow M_n$. A map $\langle f, \iota_f \rangle$ is called *co-Cartesian* if ι_f is an isomorphism. For the details of this construction, in particular, for the definition of the composition of morphisms, we refer the reader to [Gr].

The category $\mathcal{C}_\#$ comes equipped with a natural forgetful projection $\tau : \mathcal{C}_\# \rightarrow \Lambda$, and this projection is a *cofibration* in the sense of [Gr]. A *section* of this projection is a functor $\sigma : \Lambda \rightarrow \mathcal{C}_\#$ such that $\tau \circ \sigma = \text{id}$ (since Λ is small, there is no harm in requiring that two functors from Λ to itself be equal, not just isomorphic). These sections obviously form a category which we denote by $\text{Sec}(\mathcal{C}_\#)$. Explicitly, an object $M_\# \in \text{Sec}(\mathcal{C}_\#)$ is given by the following:

- (i) a collection of objects $M_n = M_\#([n]) \in \mathcal{C}^n$, and
- (ii) a collection of transition maps $\iota_f : f_!M_{n'} \rightarrow M_n$ for any n, n' , and $f \in \Lambda([n'], [n])$,

subject to natural compatibility conditions.

A section $\sigma : \Lambda \rightarrow \mathcal{C}_\#$ is called *co-Cartesian* if $\sigma(f)$ is a cocartesian map for any $[n], [n'] \in \Lambda$ and $f : [n'] \rightarrow [n]$, equivalently, a section is cocartesian if all the transition maps ι_f are isomorphisms. co-Cartesian sections form a full subcategory $\text{Sec}_{\text{cart}}(\mathcal{C}_\#)$.

Lemma 2.2. *The category $\mathbf{Sec}_{\text{cart}}(\mathcal{C}_{\#})$ of co-Cartesian objects $M_{\#} \in \mathbf{Sec}(\mathcal{C}_{\#})$ is equivalent to the category of the following data:*

- (i) an object $M = M_{\#}([1]) \in \mathcal{C}$, and
 - (ii) an isomorphism $\tau : I \times M \rightarrow M \times I$ in the category $\mathcal{C}^2 = \mathcal{C} \times \mathcal{C}$,
- such that if we denote by τ_{ij} the endomorphism of $I \times I \times M \in \mathcal{C}^3$ obtained by applying τ to the i -th and j -th multiples, we have $\tau_{31} \circ \tau_{12} \circ \tau_{23} = \text{id}$.

Proof. Straightforward and left to the reader.

Thus the natural forgetful functor $\mathbf{Sec}_{\text{cart}}(\mathcal{C}_{\#}) \rightarrow \mathcal{C}$, $M_{\#} \mapsto M_{\#}([1])$ is faithful: an object in $\mathbf{Sec}_{\text{cart}}(\mathcal{C}_{\#})$ is given by $M_{\#}([1])$ plus some extra structure on it, and all the higher components $M_{\#}([n])$, $n \geq 2$, together with the transition maps ι_f , can be recovered from $M_{\#}([1])$ and this extra structure.

Return now to the abelian situation: we are given an associative unital algebra A over a field k , and our monoidal category is $\mathcal{C} = A\text{-bimod}$, with the natural tensor product. Then for every n , the product $A\text{-bimod}^n$ has a fully faithful embedding $A\text{-bimod}^n \rightarrow A^{\otimes n}\text{-bimod}$, $M_1 \times M_2 \times \cdots \times M_n \mapsto M_1 \boxtimes M_2 \boxtimes \cdots \boxtimes M_n$, and one checks easily that the multiplication functors m_S actually extend to right-exact functors

$$m_S : A^{\otimes S}\text{-bimod} \rightarrow A\text{-bimod};$$

for instance, one can define m_S as

$$m_S(M) = M / \{a_{v'}m - ma_v \mid v \in S, a \in A, m \in M\},$$

where $a_v = 1 \otimes \cdots \otimes a \otimes \cdots \otimes 1 \in A^{\otimes S}$ with a at the v -th position, and $v' \in S$ is the next element after v . We can therefore define the cofibered category $A\text{-bimod}_{\#}/A$ with fiber $A^{\otimes V([n])}\text{-bimod}$ over $[n] \in A$, and transition functors f_i as in (2.5). We also have the category of sections $\mathbf{Sec}(A\text{-bimod}_{\#})$ and the subcategory of co-Cartesian sections $\mathbf{Sec}_{\text{cart}}(A\text{-bimod}_{\#}) \subset \mathbf{Sec}(A\text{-bimod}_{\#})$.

Lemma 2.3. *The category $\mathbf{Sec}(A\text{-bimod}_{\#})$ is a k -linear abelian category.*

Proof (Sketch of a proof). This is a general fact about cofibered categories; the proof is straightforward. The kernel $\mathbf{Ker} \phi$ and cokernel $\mathbf{Coker} \phi$ of a map $\phi : M_{\#} \rightarrow M'_{\#}$ between objects $M_{\#}, M'_{\#} \in \mathbf{Sec}(A\text{-bimod}_{\#})$ are taken pointwise: for every n , we have an exact sequence

$$0 \rightarrow (\mathbf{Ker} \phi)([n]) \rightarrow M_{\#}([n]) \xrightarrow{\phi} M'_{\#}([n]) \rightarrow (\mathbf{Coker} \phi)([n]) \rightarrow 0.$$

The transition maps ι_f for $\mathbf{Ker} \phi$ are obtained by restriction from those for $M_{\#}$; for $\mathbf{Coker} \phi$, one uses the fact that the functors f_i are right-exact.

Definition 2.4. *A cyclic bimodule M over a unital associative algebra A is a cocartesian section $M_{\#} \in \mathbf{Sec}_{\text{cart}}(A\text{-bimod}_{\#})$. A complex of cyclic bimodules M_{\bullet} over A is an object in the derived category $\mathcal{D}(\mathbf{Sec}(A\text{-bimod}_{\#}))$ whose homology objects are co-Cartesian.*

Complexes of cyclic bimodules obviously form a full triangulated subcategory in $\mathcal{D}(\text{Sec}(A\text{-bimod}_{\#}))$; consistent notation for the category would be $\mathcal{D}_{\text{cart}}(\text{Sec}(A\text{-bimod}_{\#}))$, but for simplicity we will denote its $\mathcal{DA}(A\text{-bimod})$. We have to define complexes separately for the following reasons:

- (i) The category $\text{Sec}_{\text{cart}}(A\text{-bimod}_{\#}) \subset \text{Sec}(A\text{-bimod}_{\#})$ need not be abelian: since the transition functors f_i are only right-exact, the condition of being cocartesian need not be preserved in passing to kernels.
- (ii) Even if $\text{Sec}_{\text{cart}}(A\text{-bimod}_{\#})$ is abelian, its derived category might be much smaller than $\mathcal{DA}(A\text{-bimod})$.

Example 2.5. An extreme example of (ii) is the case $A = k$: in this case $\text{Sec}(A\text{-bimod}_{\#})$ is just the category of cyclic vector spaces, $\text{Fun}(A, k)$, and $E \in \text{Fun}(A, k)$ is co-Cartesian if and only if $E(f)$ is invertible for any map $f : [n'] \rightarrow [n]$. One deduces easily that E must be a constant functor, so that $\text{Sec}_{\text{cart}}(k\text{-bimod}_{\#}) = k\text{-Vect}$. Then $\mathcal{DA}(k\text{-bimod})$ is the full subcategory $\mathcal{D}_{\text{const}}(A, k) \subset \mathcal{D}(A, k)$ of complexes whose homology is constant. If we were to consider Δ^{opp} instead of A , we would have $\mathcal{D}_{\text{const}}(\Delta^{\text{opp}}, k) \cong \mathcal{D}(k\text{-Vect})$: since $H^{\bullet}(\Delta^{\text{opp}}, k) = k$, the embedding $\mathcal{D}(k\text{-Vect}) \rightarrow \mathcal{D}(\Delta^{\text{opp}}, k)$ is fully faithful, and $\mathcal{D}_{\text{const}}(\Delta^{\text{opp}}, k)$ is its essential image. However, $H^{\bullet}(A, k)$ is $k[u]$, not k . Therefore there are maps between constant functors in $\mathcal{D}(A, k)$ that do not come from maps in $\mathcal{D}(k\text{-Vect})$, and the cones of these maps give objects in $\mathcal{D}_{\text{const}}(A, k)$ that do not come from $\mathcal{D}(k\text{-Vect})$.

This phenomenon is quite common in homological algebra; examples are, for instance, the triangulated category of complexes of étale sheaves with constructible homology, the category of complex of \mathcal{D} -modules with holonomic homology, and the so-called “equivariant derived category” of sheaves on a topological space X acted upon by a topological group G (which is not in fact the derived category of anything useful). The upshot is that it is the triangulated category $\mathcal{DA}(A\text{-bimod})$ that should be treated as the basic object, wherever categories are discussed.

Remark 2.6. We note one interesting property of the category $\mathcal{D}_{\text{const}}(A, k)$. Fix an integer $n \geq 1$, and consider the full subcategory $\Lambda_{\leq n} \subset \Lambda$ of objects $[n'] \in \Lambda$ with $n' \leq n$. Then one can show that $H^{\bullet}(\Lambda_{\leq n}, k) = k[u]/u^n$, so that we have a natural exact triangle

$$H_{\bullet}(\Lambda_{\leq n}, E^{\bullet}) \longrightarrow HC_{\bullet}(E^{\bullet}) \xrightarrow{u^n} HC_{\bullet+2n}(E) \longrightarrow, \quad (2.6)$$

for every $E^{\bullet} \in \mathcal{D}_{\text{const}}(A, k)$. We note that for any $E^{\bullet} \in \mathcal{D}(A, k)$, (1.4) extends to a spectral sequence

$$HH_{\bullet}(E^{\bullet})[u^{-1}] \Rightarrow HC_{\bullet}(E), \quad (2.7)$$

where the expression on the left-hand side reads as “polynomials in one formal variable u^{-1} of homological degree 2 with coefficients in $HH_{\bullet}(E^{\bullet})$.” Then (2.6)

shows that for $E^\bullet \in \mathcal{D}_{const}(A, k)$, the first n differentials in (2.7) depend only on the restriction of E^\bullet to $A_{\leq(n+1)} \subset A$. This is useful because in practice, one is often interested only in the first differential in the spectral sequence.

As in Lemma 2.2, a cyclic A -bimodule $M_\#$ essentially consists of an A -bimodule $M = M_\#([1])$ equipped with an extra structure. Explicitly, this structure is a map $\tau : A \otimes_k M \rightarrow M \otimes_k A$ that respects the $A^{\otimes 2}$ -bimodule structure on both sides and satisfies the condition $\tau_{31} \circ \tau_{12} \circ \tau_{23} = \text{id}$, as in Lemma 2.2.

Another way to view this structure is the following. One checks easily that for any cyclic A -bimodule $M_\#$, the restriction $j^*M_\# \in \text{Fun}(\Delta^{opp}, k)$ is canonically isomorphic to the simplicial k -vector space $M_\#^\Delta$ associated to the underlying A -bimodule M as in (2.3). By adjunction, we have a natural map

$$\tau_\# : j_!M_\#^\Delta \rightarrow M_\#.$$

Then $j_!M_\#^\Delta$ in this formula depends only on $M \in A\text{-bimod}$, and all the structure maps that turn M into the cyclic bimodule $M_\#$ are collected in the map $\tau_\#$.

We can now define cyclic homology with coefficients. The definition is rather tautological. We note that for any cyclic A -bimodule $M_\#$, or in fact, for any $M_\# \in \text{Sec}(A\text{-bimod}_\#)$, we can treat $M_\#$ as a cyclic vector space by forgetting the bimodule structure on its components M_n .

Definition 2.7. *The cyclic homology $HC_\bullet(A, M_\#)$ with coefficients in a cyclic A -bimodule M is equal to $H_\bullet(A, M_\#)$.*

Of course, (1.4), being valid for any cyclic k -vector space, also applies to $HC_\bullet(A, M_\#)$, so that we automatically get the whole package: the Connes exact sequence, the periodicity endomorphism, and the periodic cyclic homology $HP_\bullet(A, M)$. By Lemma 2.1, $HH_\bullet(M_\#)$ coincides with $HH_\bullet(A, M)$ as defined in (2.2).

3 Gauss–Manin connection.

To illustrate the usefulness of the notion of a cyclic bimodule, let us study the behavior of cyclic homology under deformations.

There are two types of deformation theory objects that one can study for an associative algebra A . The first is the notion of a *square-zero extension* of the algebra A by an A -bimodule M . This is an associative algebra \tilde{A} that fits into a short exact sequence

$$0 \longrightarrow M \xrightarrow{i} \tilde{A} \xrightarrow{p} A \longrightarrow 0,$$

where p is an algebra map, and i is an \tilde{A} -bimodule map, under the \tilde{A} -bimodule structure on M induced from the given A -bimodule structure by means of the

map p . In other words, the multiplication on the ideal $\text{Ker } p \subset \tilde{A}$ is trivial, so that the \tilde{A} -bimodule structure on $\text{Ker } p$ is induced by an A -bimodule structure, and i identifies the A -bimodule $\text{Ker } p$ with M . Square-zero extensions are classified up to an isomorphism by elements in the second Hochschild cohomology group $HH^2(A, M)$, defined as

$$HH^\bullet(A, M) = \text{Ext}_{A^{opp} \otimes A}^\bullet(A, M).$$

In this setting, we can consider the cyclic homology of the algebra \tilde{A} and compare it with the cyclic homology of A . T. Goodwillie's theorem [Go] claims that if the base field k has characteristic 0, the natural map

$$HP_\bullet(\tilde{A}) \rightarrow HP_\bullet(A)$$

is an isomorphism, and there is also some information on the behavior of $HC_\bullet(A)$.

A second type of deformation-theory data includes a commutative k -algebra R with a maximal ideal $\mathfrak{m} \subset R$. A *deformation* A_R of the algebra A over R is a flat associative unital algebra A_R over R equipped with an isomorphism $A_R/\mathfrak{m} \cong A$. In this case, one can form the *relative cyclic R -module* $A_{R\#}$ by taking the tensor products over R ; thus we have relative homology $HH_\bullet(A_R/R)$, $HC_\bullet(A_R/R)$, $HP_\bullet(A_R/R)$. The fundamental fact discovered by E. Getzler [Ge] is that we have an analogue of the Gauss–Manin connection: if $\text{Spec}(R)$ is smooth, the R -module $HP_i(A_R/R)$ carries a canonical flat connection for every i .

Consider now the case that R is not smooth but, on the contrary, local Artin. Moreover, assume that $\mathfrak{m}^2 = 0$, so that R is itself a (commutative) square-zero extension of k . Then a deformation A_R of A over R is also a square-zero extension of A , by the bimodule $A \otimes \mathfrak{m}$ (\mathfrak{m} here is taken as a k -vector space). But this square-zero extension is special: for a general square-zero extension \tilde{A} of A by some $M \in A\text{-bimod}$, there does not exist any analogue of the relative cyclic R -module $A_{R\#} \in \text{Fun}(A, R)$.

We observe the following: the data needed to define such an analogue is precisely a cyclic bimodule structure on the bimodule M .

Namely, assume given a square-zero extension \tilde{A} of the algebra A by some A -bimodule M , and consider the cyclic k -vector space $\tilde{A}_\# \in \text{Fun}(A, k)$. Let us equip \tilde{A} with a decreasing two-step filtration F^\bullet by setting $F^1 \tilde{A} = \tilde{M}$. Then this induces a decreasing filtration F^\bullet on tensor powers $\tilde{A}^{\otimes n}$. Since \tilde{A} is square-zero, F^\bullet is compatible with the multiplication maps; therefore we also have a filtration F^\bullet on $\tilde{A}_\#$. Consider the quotient

$$\overline{A_\#} = \tilde{A}_\# / F^2 \tilde{A}_\#.$$

One checks easily that $\text{gr}_F^0 \tilde{A}_\# \cong A_\#$ and $\text{gr}_F^1 \tilde{A}_\# \cong j_! M_\#^\Delta$ in a canonical way, so that $\overline{A_\#}$ fits into a canonical short exact sequence

$$0 \longrightarrow j_! M_{\#}^{\Delta} \longrightarrow \overline{A_{\#}} \longrightarrow A_{\#} \longrightarrow 0 \quad (3.1)$$

of cyclic k -vector spaces.

Now assume in addition that M is equipped with a structure of a cyclic A -bimodule $M_{\#}$, so that $M_{\#}^{\Delta} \cong j^* M_{\#}$, and we have the structure map $\tau_{\#} : j_! M_{\#}^{\Delta} \rightarrow M_{\#}$. Then we can compose the extension (3.1) with the map $\tau_{\#}$, to obtain a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & j_! M_{\#}^{\Delta} & \longrightarrow & \overline{A_{\#}} & \longrightarrow & A_{\#} \longrightarrow 0 \\ & & \tau_{\#} \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & M_{\#} & \longrightarrow & \widehat{A_{\#}} & \longrightarrow & A_{\#} \longrightarrow 0 \end{array} \quad (3.2)$$

of short exact sequences in $\text{Fun}(A, k)$, with Cartesian left square. It is easy to check that when $\tilde{A} = A_R$ for some square-zero R , so that $M = A \otimes \mathfrak{m}$, and we take the cyclic A -bimodule structure on M induced by the tautological structure on A , then $\widehat{A_{\#}}$ coincides precisely with the relative cyclic object $A_{R\#}$ (which we consider as a k -vector space, forgetting the R -module structure).

We believe that this is the proper generality for the Getzler connection; in this setting, the main result reads as follows.

Proposition 3.1. *Assume given a square-zero extension \tilde{A} of an associative algebra A by an A -bimodule M , and assume that M is equipped with a structure of a cyclic A -bimodule. Then the long exact sequence*

$$HP_{\bullet}(A, M) \longrightarrow HP_{\bullet}(\widehat{A_{\#}}) \longrightarrow HP_{\bullet}(A) \longrightarrow$$

of periodic cyclic homology induced by the second row in (3.2) admits a canonical splitting $HP_{\bullet}(A) \rightarrow HP_{\bullet}(\widehat{A_{\#}})$.

Proof. By definition, we have two natural maps

$$\begin{aligned} HP_{\bullet}(\overline{A_{\#}}) &\rightarrow HP_{\bullet}(A_{\#}) = HP_{\bullet}(A), \\ HP_{\bullet}(\overline{A_{\#}}) &\rightarrow HP_{\bullet}(\widehat{A_{\#}}), \end{aligned} \quad (3.3)$$

and the cone of the first map is isomorphic to $HP_{\bullet}(j_! M_{\#}^{\Delta})$. Since $j_!$ is exact, we have $HC_{\bullet}(j_! M_{\#}^{\Delta}) \cong HH_{\bullet}(M_{\#})$, and the periodicity map $u : HC_{\bullet}(j_! M_{\#}^{\Delta}) \rightarrow HC_{\bullet-2}(j_! M_{\#}^{\Delta})$ is equal to 0, so that $HP_{\bullet}(j_! M_{\#}^{\Delta}) = 0$. Thus the first map in (3.3) is an isomorphism, and the second map is then the required splitting.

Corollary 3.2. *Assume given a commutative k -algebra R with a maximal ideal $\mathfrak{m} \subset R$, and a deformation A_R of the algebra A over R . Then if $\text{Spec}(R)$ is smooth, the R -modules $HP_{\bullet}(A_R/R)$ carry a natural connection.*

Proof (Sketch of a proof). Consider the $R \otimes R$ -algebras $A_R \otimes R$ and $R \otimes A_R$, and their restrictions to the first infinitesimal neighborhood of the diagonal in $\mathrm{Spec}(R \otimes R) = \mathrm{Spec}(R) \times \mathrm{Spec}(R)$. Then Proposition 3.1, suitably generalized, shows that $HP_\bullet(-)$ of these two restrictions are canonically isomorphic. It is well known that giving such an isomorphism is equivalent to giving a connection on $HP_\bullet(A_R/R)$.

We note that we do not claim that the connection is *flat*. It certainly is, at least in characteristic 0; but our present method does not allow one to go beyond square-zero extensions. Thus we cannot analyze the second infinitesimal neighborhood of the diagonal in $\mathrm{Spec}(R \otimes R)$, and we cannot prove flatness.

Unfortunately, at present, we do not understand what is the proper cyclic bimodule context for higher-level infinitesimal extensions. Of course, if one is interested only in an R -deformation $\tilde{A} = A_R$ over an Artin local base R , not in its cyclic bimodule generalizations, one can use Goodwillie's Theorem: using the full cyclic object $\tilde{A}_\#$ instead of its quotient $\overline{\tilde{A}_\#}$ in Proposition 3.1 immediately gives a splitting $HP_\bullet(A) \rightarrow HP_\bullet(A_R/R)$ of the augmentation map $HP_\bullet(A_R/R) \rightarrow HP_\bullet(A)$, and this extends by R -linearity to an isomorphism $HP_\bullet(A_R/R) \cong HP_\bullet(A) \otimes R$. However, this is not quite satisfactory from the conceptual point of view, and it does not work in positive characteristic (where Goodwillie's Theorem is simply not true). If $\mathrm{char} k \neq 2$, the latter can be cured by using $\tilde{A}_\#/F^3\tilde{A}_\#$, but the former one remains. We plan to return to this elsewhere.

4 Categorical Approach.

Let us now try to define cyclic homology in a more general setting: we will attempt to replace A -bimod with an arbitrary associative unital k -linear tensor category \mathcal{C} with a unit object $\mathbf{1} \in \mathcal{C}$. We do not assume that \mathcal{C} is symmetric in any way. However, we will assume that the tensor product $- \otimes -$ is right-exact in each variable, and we will need to impose additional technical assumptions later on.

The first thing to do is to try to define Hochschild homology; so let us look more closely at (2.1). The formula on the right-hand side looks symmetric, but this is an optical illusion; the two copies of A are completely different objects: one is a left module over $A^{opp} \otimes A$, and the other is a right module (A just happens to have both structures at the same time). It is better to separate them and introduce the functor

$$\mathrm{tr} : A\text{-bimod} \rightarrow k\text{-Vect}$$

by $\mathrm{tr}(M) = M \otimes_{A^{opp} \otimes A} A$, or equivalently, by

$$\mathrm{tr}(M) = M / \{am - ma \mid a \in A, m \in M\}. \quad (4.1)$$

Then tr is a right-exact functor, and we have $HH_\bullet(A, M) = L^\bullet \mathrm{tr}(M)$.

We want to emphasize that the functor tr cannot be recovered from the tensor structure on $A\text{-bimod}$; this really is an extra piece of data. For a general tensor category \mathcal{C} , it does not exist a priori; we have to impose it as an additional structure.

Let us axiomatize the situation. First, forget for the moment about the k -linear and abelian structure on \mathcal{C} ; let us treat it simply as a monoidal category. Assume given some other category \mathcal{B} and a functor $T : \mathcal{C} \rightarrow \mathcal{B}$.

Definition 4.1. *The functor $T : \mathcal{C} \rightarrow \mathcal{B}$ is a trace functor if it is extended to a functor $\mathcal{C}_\# \rightarrow \mathcal{B}$ that sends any cocartesian map $f : M \rightarrow M'$ in $\mathcal{C}_\#$ to an invertible map.*

Another way to say the same thing is the following: the categories $\text{Fun}(\mathcal{C}^n, \mathcal{B})$ of functors from \mathcal{C}^n to \mathcal{B} form a fibered category over Λ , and a trace functor is a Cartesian section of this fibration. Explicitly, a trace functor is defined by $T : \mathcal{C} \rightarrow \mathcal{B}$ and a collection of isomorphisms

$$T(M \otimes M') \rightarrow T(M' \otimes M)$$

for any $M, M' \in \mathcal{C}$ that are functorial in M and M' and satisfy some compatibility conditions analogous to those in Lemma 2.2; we leave it to the reader to write down these conditions precisely. Thus T has a tracelike property with respect to the product in \mathcal{C} , and this motivates our terminology.

Recall now that \mathcal{C} is a k -linear abelian category. To define Hochschild homology, we have to assume that it is equipped with a right-exact trace functor $\text{tr} : \mathcal{C} \rightarrow k\text{-Vect}$; then for any $M \in \mathcal{C}$, we set

$$HH_*(M) = L^\bullet \text{tr}(M). \tag{4.2}$$

Lemma 4.2. *The functor $\text{tr} : A\text{-bimod} \rightarrow k\text{-Vect}$ canonically extends to a right-exact trace functor in the sense of Definition 4.1.*

Proof. For any object $\langle [n], M_n \rangle \in A\text{-bimod}_\#$, $[n] \in \Lambda$, $M_n \in A^{\otimes n}\text{-bimod}$, let

$$\text{tr}(\langle [n], M_n \rangle) = M_n / \{ a_{v'} m - m a_v \mid v \in V([n]), m \in M_n, a \in A \},$$

where $a_v = 1 \otimes 1 \otimes \dots \otimes a \otimes \dots \otimes 1 \in A^{\otimes V([n])}$ has a in the multiple corresponding to $v \in V([n])$, and $v' \in V([n])$ is the next marked point after v counting clockwise. The compatibility with maps in the category $A\text{-bimod}_\#$ is obvious.

We note that here, in the case $\mathcal{C} = A\text{-bimod}$, the category $A\text{-bimod}_\#$ is actually larger than what we would have had purely from the monoidal structure on \mathcal{C} : M_n is allowed to be an arbitrary $A^{\otimes n}$ -bimodule, not a collection of n A -bimodules. To do the same for general k -linear \mathcal{C} , we need to replace $A^{\otimes n}\text{-bimod}$ with some version of the tensor product $\mathcal{C}^{\otimes n}$. Here we have a difficulty: for various technical reasons, it is not clear how to define tensor products for sufficiently general abelian categories.

One way around it is the following. For any (small) k -linear abelian category \mathcal{B} , a k -linear functor $\mathcal{B}^{opp} \rightarrow k\text{-Vect}$ is left-exact if and only if it is a sheaf for the canonical Grothendieck topology on \mathcal{B} ([BD, 5, §10]); the category $\text{Shv}(\mathcal{B})$ of such functors is abelian and k -linear, and \mathcal{B} itself is naturally embedded into $\text{Shv}(\mathcal{B})$ by Yoneda. The embedding is a fully faithful exact functor. Every functor in $\text{Shv}(\mathcal{B})$ is in fact a direct limit of representable functors, so that $\text{Shv}(\mathcal{B})$ is an inductive completion of the abelian category \mathcal{B} . Now, if we are given two (small) k -linear abelian categories $\mathcal{B}_1, \mathcal{B}_2$, then their product $\mathcal{B}_1 \times \mathcal{B}_2$ is no longer abelian. However, we still have the abelian category $\text{Shv}(\mathcal{B}_1 \times \mathcal{B}_2)$ of bilinear functors $\mathcal{B}_1^{opp} \times \mathcal{B}_2^{opp} \rightarrow k\text{-Vect}$, which are left-exact in each variable, and the same goes for polylinear functors.

Moreover, for any right-exact functor $F : \mathcal{B}_1 \rightarrow \mathcal{B}_2$ between small abelian categories, we have the restriction functor $F^* : \text{Shv}(\mathcal{B}_2) \rightarrow \text{Shv}(\mathcal{B}_1)$, which is left-exact, and its left-adjoint $F_! : \text{Shv}(\mathcal{B}_1) \rightarrow \text{Shv}(\mathcal{B}_2)$, which is right-exact. The functor $F_!$ is an extension of the functor F : on Yoneda images $\mathcal{B}_i \subset \text{Shv}(\mathcal{B}_i)$, we have $F_! = F$. And again, the same works for polylinear functors.

In particular, given our k -linear abelian tensor category \mathcal{C} , we can form the category $\text{Shv}(\mathcal{C})_{\#}$ of pairs $\langle E, [n] \rangle$, $[n] \in \Lambda$, $E \in \text{Shv}(\mathcal{C}^n)$, with a map from $\langle E', [n'] \rangle$ to $\langle E, [n] \rangle$ given by a pair of a map $f : [n'] \rightarrow [n]$ and either a map $E' \rightarrow (f_!)^* E$, or map $(f_!)_! E' \rightarrow E$; this is equivalent by adjunction. Then $\text{Shv}(\mathcal{C})_{\#}$ is a bifibered category over Λ in the sense of [Gr].

The category of sections $\Lambda \rightarrow \text{Shv}(\mathcal{C})_{\#}$ of this bifibration can also be described as the full subcategory $\text{Shv}(\mathcal{C}_{\#}) \subset \text{Fun}(\mathcal{C}_{\#}^{opp}, k)$ spanned by those functors $E_{\#} : \mathcal{C}_{\#}^{opp} \rightarrow k\text{-Vect}$ whose restriction to $(\mathcal{C}^{opp})^n \subset \mathcal{C}_{\#}^{opp}$ is a sheaf, that is, an object in $\text{Shv}(\mathcal{C}^n) \subset \text{Fun}((\mathcal{C}^{opp})^n, k)$. Since the transition functors $(f_!)_!$ are right-exact, $\text{Shv}(\mathcal{C}_{\#})$ is an abelian category (this is proved in exactly the same way as Lemma 2.3).

We denote by $\text{Shv}_{cart}(\mathcal{C}_{\#}) \subset \text{Shv}(\mathcal{C}_{\#})$ the full subcategory of sections $E : \Lambda \rightarrow \text{Shv}(\mathcal{C})_{\#}$ that are co-Cartesian, and moreover, are such that $E([1]) \in \text{Shv}(\mathcal{C})$ actually lies in the Yoneda image $\mathcal{C} \subset \text{Shv}(\mathcal{C})$. We also denote by $\mathcal{DA}(\mathcal{C}) \subset \mathcal{D}(\text{Shv}(\mathcal{C}_{\#}))$ the full triangulated subcategory of complexes $E_{\#}^{\bullet} \in \mathcal{D}(\text{Shv}(\mathcal{C}_{\#}))$ with homology in $\text{Shv}_{cart}(\mathcal{C}_{\#})$.

If \mathcal{C} is the category of A -bimodules for some algebra A , or better yet, of A -bimodules of cardinality not more than that of $A \times \mathbb{N}$, so that \mathcal{C} is small, then $\text{Shv}(\mathcal{C})$ is equivalent to $A\text{-bimod}$ (one shows easily that every sheaf $E \in \text{Shv}(\mathcal{C})$ is completely determined by its value at $A^{opp} \otimes A \in \mathcal{C}$). In this case, $\mathcal{DA}(\mathcal{C})$ is our old category $\mathcal{DA}(A\text{-bimod})$.

Now we assume that \mathcal{C} is equipped with a right-exact trace functor $\text{tr} : \mathcal{C} \rightarrow k\text{-Vect}$. We would like to define cyclic homology $HC_{\bullet}(M_{\bullet})$ for any $M_{\bullet} \in \mathcal{DA}(\mathcal{C})$, and we immediately notice a problem: for a general \mathcal{C} , we do not have a forgetful functor to vector spaces. However, it turns out that the forgetful functor is *not needed* for the definition; it can be replaced with the trace functor tr .

We proceed as follows. By definition, tr is extended to a functor $\mathcal{C}_{\#} \rightarrow k\text{-Vect}$; we extend it canonically to a functor $\text{Shv}(\mathcal{C})_{\#} \rightarrow k\text{-Vect}$, and consider

the product

$$\mathrm{tr} \times \tau : \mathrm{Shv}(\mathcal{C})_{\#} \rightarrow k\text{-Vect} \times \Lambda,$$

where $\tau : \mathrm{Shv}(\mathcal{C})_{\#} \rightarrow \Lambda$ is the projection. This is a functor compatible with the projections to Λ , and therefore, it induces a functor of the categories of sections. The category of sections of the projection $k\text{-Vect} \times \Lambda \rightarrow \Lambda$ is tautologically the same as $\mathrm{Fun}(\Lambda, k\text{-Vect})$, so that we have a functor

$$\mathrm{tr}_{\#} : \mathrm{Shv}(\mathcal{C}_{\#}) \rightarrow \mathrm{Fun}(\Lambda, k).$$

One checks easily that this functor is right-exact.

Definition 4.3. For any $M_{\#} \in \mathrm{Sec}(\mathcal{C}_{\#})$, its cyclic homology $HC_{\bullet}(M_{\#})$ is defined by

$$HC_{\bullet}(M_{\#}) \stackrel{\mathrm{def}}{=} HC_{\bullet}(L^{\bullet} \mathrm{tr}_{\#}(M_{\#})) = H_{\bullet}(\Lambda, L^{\bullet} \mathrm{tr}_{\#}(M_{\#})).$$

Definition 4.4. The pair $\langle \mathcal{C}, \mathrm{tr} \rangle$ is called homologically clean if for any n , the category $\mathrm{Shv}(\mathcal{C}^n)$ has enough objects E such that

- (i) E is acyclic both for functors $(f_!)_! : \mathrm{Shv}(\mathcal{C}^n) \rightarrow \mathrm{Shv}(\mathcal{C}^{n'})$, for any $f : [n] \rightarrow [n']$, and for the trace functor $\mathrm{tr} : \mathrm{Shv}(\mathcal{C}^n) \rightarrow k\text{-Vect}$, and
- (ii) for any $f : [n] \rightarrow [n']$, $(f_!)_! E \in \mathrm{Shv}(\mathcal{C}^{n'})$ is acyclic for $\mathrm{tr} : \mathrm{Shv}(\mathcal{C}^{n'}) \rightarrow k\text{-Vect}$.

Example 4.5. Assume that the category \mathcal{C} has enough projectives, and moreover, $P_1 \otimes P_2$ is projective for any projective $P_1, P_2 \in \mathcal{C}$ (this is satisfied, for instance, for $\mathcal{C} = A\text{-bimod}$). Then the pair $\langle \mathcal{C}, \mathrm{tr} \rangle$ is homologically clean, for any trace functor tr . Indeed, $\mathrm{Shv}(\mathcal{C}^n)$ then also has enough projectives, say sums of objects

$$P = P_1 \boxtimes P_2 \boxtimes \cdots \boxtimes P_n \in \mathrm{Shv}(\mathcal{C}^n) \tag{4.3}$$

for projective $P_1, \dots, P_n \in \mathcal{C} \subset \mathrm{Shv}(\mathcal{C})$, and these projectives automatically satisfy the condition (i). To check (ii), one decomposes $f : [n] \rightarrow [n']$ into a surjection $p : [n] \rightarrow [n'']$ and an injection $i : [n''] \rightarrow [n']$. Since the tensor product of projective objects is projective, $(p_!)_!(P) \in \mathrm{Shv}(\mathcal{C}^{n''})$ is also an object of type (4.3), so we may as well assume that f is injective. Then one can find a left-inverse map $f' : [n'] \rightarrow [n]$, $f' \circ f = \mathrm{id}$; since $P' = (f_!)_!(P)$ is obviously acyclic for $(f'_!)_!$, and $(f'_!)_!(P') = ((f' \circ f)_!)_!(P) = P$ is acyclic for tr , P' itself is acyclic for $\mathrm{tr} = \mathrm{tr} \circ (f'_!)_!$.

Lemma 4.6. Assume that $\langle \mathcal{C}, \mathrm{tr} \rangle$ is homologically clean. Then for any object $[n] \in \Lambda$ and any $M_{\#} \in \mathrm{Shv}(\mathcal{C}_{\#})$, we have

$$L^{\bullet} \mathrm{tr}_{\#}(M_{\#})([n]) \cong L^{\bullet} \mathrm{tr}(M_{\#}([n])). \tag{4.4}$$

For any $M_{\#} \in \mathcal{D}\Lambda(\mathcal{C})$, we have $L^{\bullet} \mathrm{tr}_{\#}(M_{\#}) \in \mathcal{D}_{\mathrm{const}}(\Lambda, k) \subset \mathcal{D}(\Lambda, k)$.

Proof. The natural restriction functor $\mathrm{Shv}(\mathcal{C}_\#) \rightarrow \mathrm{Shv}(\mathcal{C}^n)$, $M_\# \mapsto M_\#([m])$ has a left-adjoint functor $I_{n!} : \mathrm{Shv}(\mathcal{C}^n) \rightarrow \mathrm{Shv}(\mathcal{C}_\#)$; explicitly, it is given by

$$I_{n!}(E)([n']) = \bigoplus_{f:[n] \rightarrow [n']} (f!)_!(E). \quad (4.5)$$

Let us say that an object $E \in \mathrm{Shv}(\mathcal{C}^n)$ is admissible if it satisfies the conditions (i), (ii) of Definition 4.4. By assumption, $\mathrm{Shv}(\mathcal{C}^n)$ has enough admissible objects for any n . Then $\mathrm{Shv}(\mathcal{C}_\#)$ has enough objects of the form $I_{n!}E$, $[n] \in \Lambda$, $E \in \mathrm{Shv}(\mathcal{C}^n)$ admissible, and to prove the first claim, it suffices to consider $M_\# = I_{n!}E$ of this form. In degree 0, (4.4) is the definition of the functor $\mathrm{tr}_\#$, and the higher-degree terms on the right-hand side vanish by Definition 4.4 (ii). Therefore it suffices to prove that $M_\# = I_{n!}E$ is acyclic for the functor $\mathrm{tr}_\#$. This is obvious: applying $\mathrm{tr}_\#$ to any short exact sequence

$$0 \longrightarrow M'_\# \longrightarrow M''_\# \longrightarrow M_\# \longrightarrow 0$$

in $\mathrm{Shv}(\mathcal{C}_\#)$, we see that since $M_\#[[n']]$ is acyclic for any $[n'] \in \Lambda$, the sequence

$$0 \longrightarrow \mathrm{tr} M'_\#[[n']] \longrightarrow \mathrm{tr} M''_\#[[n']] \longrightarrow \mathrm{tr} M_\#[[n']] \longrightarrow 0$$

is exact; this means that

$$0 \longrightarrow \mathrm{tr} M'_\# \longrightarrow \mathrm{tr} M''_\# \longrightarrow \mathrm{tr} M_\# \longrightarrow 0$$

is an exact sequence in $\mathrm{Fun}(\Lambda, k)$, and this means that $M_\#$ is indeed acyclic for $\mathrm{tr}_\#$.

With the first claim proved, the second amounts to showing that the natural map

$$L^\bullet \mathrm{tr} \circ L^\bullet (f!)_!(E) \rightarrow L^\bullet \mathrm{tr}(E)$$

is a quasi-isomorphism for any $f : [n] \rightarrow [n']$ and any $E \in \mathrm{Shv}(\mathcal{C}^n)$. It suffices to prove it for admissible M ; then the higher derived functors vanish, and the isomorphism $\mathrm{tr} \circ (f!)_! \cong \mathrm{tr}$ is Definition 4.1.

Lemma 4.7. *In the assumptions of Lemma 4.6, for any complex $M_\#^\bullet \in \mathcal{D}\Lambda(\mathcal{C})$ with the first component $M^\bullet = M_\#^\bullet([1])$ we have*

$$HH_\bullet(M^\bullet) \cong HH_\bullet(L^\bullet \mathrm{tr}_\#(M_\#^\bullet)).$$

Proof. By Lemma 4.6, the left-hand side, $HH_\bullet(M^\bullet)$, is canonically isomorphic to the complex $L^\bullet \mathrm{tr}_\#(M_\#^\bullet) \in \mathcal{D}(\Lambda, k)$ evaluated at $[1] \in \Lambda$, and moreover, $L^\bullet \mathrm{tr}_\#(M_\#^\bullet)$ lies in the subcategory $\mathcal{D}_{\mathrm{const}}(\Lambda, k) \subset \mathcal{D}(\Lambda, k)$. It remains to apply the following general fact: for any $E^\bullet \in \mathcal{D}_{\mathrm{const}}(\Lambda, k)$, we have a natural isomorphism $HH_\bullet(E^\bullet) \cong E^\bullet([1])$. Indeed, by definition we have

$$HH_\bullet(E^\bullet) = H_\bullet(\Delta^{\mathrm{opp}}, j^* E^\bullet),$$

and $j^* E^\bullet$ lies in the category $\mathcal{D}_{\mathrm{const}}(\Delta^{\mathrm{opp}}, k)$, which is equivalent to $\mathcal{D}(k\text{-Vect})$ (see Example 2.5, and also Remark 2.6: the isomorphism we constructed here is a special case of (2.6) for $n = 1$).

The lemma shows that if the pair $\langle \mathcal{C}, \text{tr} \rangle$ is homologically clean, Definition 4.3 is consistent with (4.2), and we get the whole periodicity package of (1.4): the periodicity map u , the Connes exact sequence

$$HH_{\bullet}(M^{\bullet}) \longrightarrow HC_{\bullet}(M^{\bullet}) \xrightarrow{u} HC_{\bullet-2}(M^{\bullet}) \longrightarrow ,$$

and the periodic cyclic homology $HP_{\bullet}(M^{\bullet})$.

In general, objects in $\mathcal{DA}(\mathcal{C})$ may be hard to construct, but we always have at least one: the identity section $l_{\#} : \mathcal{A} \rightarrow \text{Shv}(\mathcal{C})_{\#}$, given by

$$l_{\#}([n]) = l^{\boxtimes n} \in \mathcal{C}^{\otimes n},$$

where $l \in \mathcal{C}$ is the unit object. Thus we can define cyclic homology of a tensor category equipped with a trace functor.

Definition 4.8. *For any k -linear abelian unital tensor category \mathcal{C} equipped with a trace functor $\text{tr} : \mathcal{C} \rightarrow k\text{Vect}$, its Hochschild and cyclic homologies are given by*

$$HH_{\bullet}(\mathcal{C}, \text{tr}) \stackrel{\text{def}}{=} HH_{\bullet}(l), \quad HC_{\bullet}(\mathcal{C}, \text{tr}) \stackrel{\text{def}}{=} HC_{\bullet}(l_{\#}),$$

where $l \in \mathcal{C}$ is the unit object, and $l_{\#} \in \mathcal{DA}(\mathcal{C})$ is the identity section.

We now have to check that in the case $\mathcal{C} = A\text{-bimod}$, Definition 4.3 is compatible with our earlier Definition 2.7, in other words, that the cyclic homology computed by means of the forgetful functor is the same as the cyclic homology computed by means of the trace. This is not at all trivial. Indeed, if for instance $M_{\#} \in \text{Shv}(\mathcal{C}_{\#})$ is co-Cartesian, then, while $L^{\bullet} \text{tr}^{\#} M_{\#}$ lies in the subcategory $\mathcal{D}_{\text{const}}(\mathcal{A}, k) \subset \mathcal{D}(\mathcal{A}, k)$, the same is certainly not true for the object $M_{\#} \in \text{Fun}(\mathcal{A}, k)$ obtained by forgetting the bimodule structure on M_n .

Thus these two objects are different. However, they do become equal after taking cyclic (or Hochschild, or periodic cyclic) homology. Namely, for any $M_{\#} \in \text{Sec}(A\text{-bimod}_{\#})$ we have a natural map

$$M_{\#} \rightarrow L^{\bullet} \text{tr}^{\#} M_{\#} \tag{4.6}$$

in the derived category $\mathcal{D}(\mathcal{A}, k)$, and we have the following result.

Proposition 4.9. *For every $M_{\#} \in \text{Sec}(A\text{-bimod}_{\#})$, the natural map (4.6) induces isomorphisms*

$$\begin{aligned} HH_{\bullet}(M_{\#}) &\cong HH_{\bullet}(L^{\bullet} \text{tr} M_{\#}), \\ HC_{\bullet}(M_{\#}) &\cong HC_{\bullet}(L^{\bullet} \text{tr} M_{\#}), \\ HP_{\bullet}(M_{\#}) &\cong HP_{\bullet}(L^{\bullet} \text{tr} M_{\#}). \end{aligned}$$

Proof. By (1.4), it suffices to consider $HC_{\bullet}(-)$; as in the proof of Lemma 4.6, it suffices to consider $M_{\#} = I_n!E$ given in (4.5), with E being the free bimodule

$$E = (A^{opp} \otimes A)^{\otimes n} \in \text{Shv}(\mathcal{C}^n) = A^{\otimes n}\text{-bimod}$$

for some fixed n . Explicitly, we have

$$I_{n!}E([n']) = \bigoplus_{f:[n] \rightarrow [n']} \bigotimes_{v' \in V([n'])} A^{opp} \otimes A^{\otimes f^{-1}(v')} \quad (4.7)$$

for any $[n'] \in \Lambda$. Then $L^p \text{tr}_{\#} I_{n!}E = 0$ for $p \geq 1$, and one checks easily that

$$\text{tr}_{\#} I_{n!}E = i_{n!} \text{tr} E = i_{n!} A^{\otimes n} \in \text{Fun}(\Lambda, k),$$

where $i_n : \text{pt} \rightarrow \Lambda$ is the embedding of the object $[n] \in \Lambda$ (pt is the category with one object and one morphism). Therefore

$$HC_0(L^{\bullet} \text{tr}_{\#} I_{n!}E) = H_{\bullet}(\Lambda, i_{n!} A^{\otimes n}) = A^{\otimes n},$$

and $HC_p(L^{\bullet} \text{tr}_{\#} i_{n!}E) = 0$ for $p \geq 1$. We have to compare it with $HC_{\bullet}(i_{n!}E)$.

To do this, consider the category $\Lambda_{[n]}$ of objects $[n'] \in \Lambda$ equipped with a map $[n] \rightarrow [n']$, and let $j_n : \Lambda_{[n]} \rightarrow \Lambda$ be the forgetful functor. Then j_n is obviously a discrete cofibration. Comparing (1.2) and (4.7), we see that

$$I_{n!}E = j_{n!}E_{\#}^{[n]}$$

for some $E_{\#}^{[n]} \in \text{Fun}(\Lambda_{[n]})$. Moreover, fix once and for all a map $[1] \rightarrow [n]$. Then we see that the discrete cofibration $j_n : \Lambda_{[n]} \rightarrow \Lambda$ factors through the discrete cofibration $j : \Lambda_{[1]} = \Delta^{opp} \rightarrow \Lambda$ by means of a discrete cofibration $\gamma_n : \Lambda_{[n]} \rightarrow \Lambda_{[1]}$, and we observe that

$$E_{\#}^{[n]}([n']) = (A^{opp})^{\otimes n'} \otimes A^{\otimes n}$$

depends only on $\gamma_n([n']) \in \Delta^{opp}$. More precisely, we have $E_{\#}^{[n]} = \gamma_n^* E_n^{\Delta}$, where $E_n^{\Delta} \in \text{Fun}(\Delta^{opp}, k)$ is as in (2.3), and E_n is the free A -bimodule

$$E_n = A^{opp} \otimes A^{\otimes(n-1)} \otimes A.$$

The conclusion: we have

$$HC_{\bullet}(I_{n!}E) = H_{\bullet}(\Lambda_{[n]}, E_{\#}^{[n]}) = H_{\bullet}(\Delta^{opp}, \gamma_{n!} \gamma_n^* E_n^{\Delta}) = H_{\bullet}(\Delta^{opp}, E_n^{\Delta} \otimes \gamma_{n!} k),$$

where we have used the projection formula (1.3) in the right-hand side. The homology of the category Δ^{opp} can be computed by the standard complex; then by the Künneth formula, the right-hand side is isomorphic to

$$H_{\bullet}(\Delta^{opp}, E_n^{\Delta}) \otimes H_{\bullet}(\Delta^{opp}, \gamma_{n!} k) \cong H_{\bullet}(\Delta^{opp}, E_n^{\Delta}) \otimes H_{\bullet}(\Lambda_{[n]}, k).$$

By Lemma 2.1,

$$H_{\bullet}(\Delta^{opp}, E_n^{\Delta}) \cong HH_{\bullet}(A, E_n) \cong A^{\otimes n}.$$

Since the category $\Lambda_{[n]}$ has an initial object $[n] \in \Lambda_{[n]}$, we have $k = i_{n!}k$, so that the second multiple $H_{\bullet}(\Lambda_{[n]}, k)$ is just k in degree 0.

The essential point of Proposition 4.9 is the following: the cyclic object $A_{\#}$ associated to an algebra A inconveniently contains two things at the same time: the cyclic structure, which seems to be essential to the problem, and the bar resolution, which is needed only to compute the Hochschild homology $HH_{\bullet}(A)$. Replacing $A_{\#}$ with the cyclic complex $L^{\bullet} \operatorname{tr}_{\#} A_{\#} \in \mathcal{D}(A, k)$ disentangles these two.

We note that while one still has to prove that this does not change the final answer, the construction itself looks pretty straightforward: if one wants to remove the inessential bar resolution from the definition of the cyclic homology, Definition 4.8 seems to be the obvious thing to try. However, it was actually arrived at by a sort of reverse engineering process. To finish the section, perhaps it would be useful to show the reader the first stage of this process.

Assume given an associative algebra A , and fix a projective resolution P_{\bullet} of the diagonal A -module A . Then $HH_{\bullet}(A, M)$ can be computed by the complex

$$\operatorname{tr}(P_{\bullet}) = P_{\bullet} \otimes_{A^{\operatorname{opp}} \otimes A} A.$$

How can one see the cyclic homology in terms of this complex? Or even simpler, what is the first differential in the spectral sequence (2.7), the Connes differential $B : HH_{\bullet}(A) \rightarrow HH_{\bullet+1}(A)$?

There is the following recipe, which gives the answer. Let $\tau : P_{\bullet} \rightarrow A$ be the augmentation map. Consider the tensor product $P_{\bullet} \otimes_A P_{\bullet}$. This is also a projective resolution of A , and we actually have *two* natural quasi-isomorphisms

$$\tau_1, \tau_2 : P_{\bullet} \otimes_A P_{\bullet} \rightarrow P_{\bullet},$$

given by $\tau_1 = \tau \otimes \operatorname{id}$, $\tau_2 = \operatorname{id} \otimes \tau$. These quasi-isomorphisms are different. However, since both are maps between projective resolutions of the same object, there should be a chain homotopy between them. Fix such a homotopy $\iota : P_{\bullet} \otimes_A P_{\bullet} \rightarrow P_{\bullet+1}$.

Now we apply the trace functor tr , and obtain two maps $\tau_1, \tau_2 :$

$\operatorname{tr}(P_{\bullet} \otimes P_{\bullet}) \rightarrow \operatorname{tr}(P_{\bullet})$, and a homotopy $\iota : \operatorname{tr}(P_{\bullet} \otimes P_{\bullet}) \rightarrow \operatorname{tr}(P_{\bullet+1})$ between them.

However, by the trace property of τ , we also have an involution $\sigma : \operatorname{tr}(P_{\bullet} \otimes_A P_{\bullet})$ that interchanges the two multiples. This involution obviously also interchanges τ_1 and τ_2 , but there is no reason why it should fix the homotopy ι – in fact, it sends ι to a second homotopy $\iota' : \operatorname{tr}(P_{\bullet} \otimes_A P_{\bullet}) \rightarrow \operatorname{tr}(P_{\bullet+1})$ between τ_1 and τ_2 .

The difference $\iota' - \iota$ is then a well-defined map of complexes

$$\iota' - \iota : \operatorname{tr}(P_{\bullet} \otimes_A P_{\bullet}) \rightarrow \operatorname{tr}(P_{\bullet+1}). \tag{4.8}$$

On the level of homology, both sides are $HH_{\bullet}(A)$; the map $\iota' - \iota$ then induces exactly the Connes differential $B : HH_{\bullet}(A) \rightarrow HH_{\bullet+1}(A)$.

To justify this recipe, we use Proposition 4.9 and identify $HC_{\bullet}(A)$ with $HC_{\bullet}(L^{\bullet} \operatorname{tr}_{\#}(A_{\#}))$ rather than $HC_{\bullet}(A_{\#})$. Then $L^{\bullet} \operatorname{tr}_{\#}(A_{\#})$ is an object in

$\mathcal{D}_{const}(\Lambda, k)$. Therefore, as noted in Remark 2.6, the Connes differential B depends only on the restriction of $L^\bullet \operatorname{tr}_\#(A_\#)$ to $\Lambda_{\leq 2} \subset \Lambda$. In other words, we do not need to compute the full $L^\bullet \operatorname{tr}_\#(A_\#)$ and to construct a full resolution $P_\#^i$ of the cyclic A -bimodule $A_\#$; it suffices to construct $P_\#^i = P_\#^{\#}([i])$ for $i = 1, 2$ (and then apply the functor tr).

With the choices made above, we set $P_\#^1 = P_\bullet$, and we let $P_\#^2$ be the cone of the map

$$P_\bullet \boxtimes P_\bullet \xrightarrow{(\tau \boxtimes \operatorname{id}) \oplus (\operatorname{id} \boxtimes \tau)} (A \boxtimes P_\bullet) \oplus (P_\bullet \boxtimes A).$$

The involution $\sigma : [2] \rightarrow [2]$ acts on $P_\#^2$ in the obvious way. We also need to define the transition maps ι_f for the two injections $d, d' : [1] \rightarrow [2]$ and the two surjections $s, s' : [2] \rightarrow [1]$. For d_1 , the transition map $\iota_d : A \boxtimes P_\bullet \rightarrow P_\#^2$ is the obvious embedding, and so is the transition map $\iota_{d'}$. For the surjection s , we need a map ι_s from the cone of the map

$$P_\bullet \otimes_A P_\bullet \xrightarrow{(\tau \otimes \operatorname{id}) \oplus (\operatorname{id} \otimes \tau)} P_\bullet \oplus P_\bullet$$

to P_\bullet . On $P_\bullet \oplus P_\bullet$, the map ι_s is just the difference map $a \oplus b \mapsto a - b$; on $P_\bullet \otimes_A P_\bullet$, ι_s is our fixed homotopy $\iota : P_\bullet \otimes_A P_\bullet \rightarrow P_{\bullet+1}$. And similarly for the other surjection s' .

We leave it to the reader to check that if one computes $L^\bullet \operatorname{tr}_\#(A_\#) |_{\Lambda_{\leq 2}}$ using this resolution $P_\#^i$, then one obtains exactly (4.8) for the Connes differential B .

5 Discussion

One of the most unpleasant features of the construction presented in Section 4 is the strong assumptions we need to impose on the tensor category \mathcal{C} . In fact, the category to which one would really like to apply the construction is the category $\operatorname{End} \mathcal{B}$ of endofunctors—whatever that means—of the category \mathcal{B} of coherent sheaves on an algebraic variety X . But if X is not affine, $\operatorname{End} \mathcal{B}$ certainly does not have enough projectives, so that Example 4.5 does not apply, and it is unlikely that $\operatorname{End} \mathcal{B}$ can be made homologically clean in the sense of Definition 4.4. We note that Definition 4.4 has been arranged so as not to impose anything more than strictly necessary for the proofs; but in practice, we do not know any examples that are not covered by Example 4.5.

As for the category $\operatorname{End} \mathcal{B}$, there is an even bigger problem with it: while there are ways to define endofunctors so that $\operatorname{End} \mathcal{B}$ is an abelian category with a right-exact tensor product, it cannot be equipped with a right-exact trace functor tr . Indeed, it immediately follows from Definition 4.8 that the Hochschild homology groups $HH_\bullet(\mathcal{C})$ of a tensor category \mathcal{C} are trivial in negative homological degrees. If $\mathcal{C} = \operatorname{End} \mathcal{B}$, one of course expects $HH_\bullet(\mathcal{C}) = HH_\bullet(X)$, the Hochschild homology $HH_\bullet(X)$ of the variety X , which by now is well understood (see, e.g., [W]). And if X is not affine, $HH_\bullet(X)$ typically is

nontrivial both in positive and in negative degrees. If X is smooth and proper, $HH_*(X)$ in fact carries a nondegenerate pairing, so that it is just as nontrivial in degrees > 0 as in degrees < 0 . Thus the case of a nonaffine algebraic variety is far beyond the methods developed in this paper.

The real reason for these difficulties is that we are dealing with abelian categories, while the theory emphatically wants to live in the triangulated world; as we explained in Example 2.5, even our main topic, cyclic bimodules, are best understood as objects of a triangulated category $\mathcal{DA}(\mathcal{C})$. Unfortunately, we cannot develop the theory from scratch in the triangulated context, since we do not have a strong and natural enough notion of an enhanced triangulated category (and working with the usual triangulated categories is out of the question because, for instance, the category of triangulated functors between triangulated categories is usually not a triangulated category itself). A well-developed theory would probably require a certain compromise between the abelian and the triangulated approaches. We will return to it elsewhere.

Another thing that is very conspicuously not done in the present paper is the combination of Section 4 and Section 3. Indeed, in Section 3, we are dealing with cyclic homology in the straightforward naive way of Section 2, and while we define the cyclic object $\widehat{A}_\#$ associated to a square-zero extension \widetilde{A} , we make no attempt to find an appropriate category $\text{Sec}(\widehat{A\text{-bimod}}_\#)$ where it should live. This is essentially the reason why we cannot go further than square-zero extensions. At present, sadly, we do not really understand this hypothetical category $\text{Sec}(\widehat{A\text{-bimod}}_\#)$.

One suspects that treating this properly would require studying deformations in a much more general context: instead of considering square-zero extensions of an algebra, we should look at the deformations of the abelian category of its modules, or at the deformations of the tensor category of its bimodules. This brings us to another topic completely untouched in the paper: the Hochschild cohomology $HH^*(A)$.

Merely *defining* Hochschild cohomology for an arbitrary tensor category \mathcal{C} is in fact much simpler than the definition of $HH_*(\mathcal{C})$, and one does not need a trace functor for this: we just set $HH^*(\mathcal{C}) = \text{Ext}^*(\mathbb{1}, \mathbb{1})$, where $\mathbb{1} \in \mathcal{C}$ is the unit object. However, it is well understood by now that just as Hochschild homology always comes equipped with the Connes differential, the spectral sequence (2.7), and the whole cyclic homology package, Hochschild cohomology should be considered not as an algebra but as the so-called *Gerstenhaber* algebra; in fact, the pair $HH_*(-), HH^*(-)$ should form a version of “noncommutative calculus,” as proposed for instance in [TT]. Deformations of the tensor category \mathcal{C} should be controlled by $HH^*(\mathcal{C})$, and the behavior of $HH_*(\mathcal{C})$ and $HC_*(\mathcal{C})$ under these deformations reflects various natural actions of $HH^*(-)$ on $HH_*(-)$.

We believe that a convenient development of the “noncommutative calculus” for a tensor category \mathcal{C} might be possible along the same lines

as our Section 4. Just as our category $\mathcal{DA}(\mathcal{C})$ is defined as the category of sections of the cofibration $\mathcal{C}_{\#}/\Lambda$, whose definition imitates the usual cyclic object $A_{\#}$, one can construct a cofibration $\mathcal{C}^{\#}/\Delta$ that imitates the standard cosimplicial object computing $HH^*(A)$: for any $[n] \in \Delta$, $\mathcal{C}^{\#}([n])$ is the category of polylinear right-exact functors from \mathcal{C}^{n-1} to \mathcal{C} , and the transition functors between various $\mathcal{C}^{\#}([n])$ are induced by the tensor product on \mathcal{C} . Then one can define a triangulated category $\mathcal{D}\Delta(\mathcal{C})$, the subcategory in $\mathcal{D}(\text{Sec}(\mathcal{C}^{\#}))$ of complexes with co-Cartesian homology; the higher structures on $HH^*(\mathcal{C})$ should be encoded in the structure of the category $\mathcal{D}\Delta(\mathcal{C})$, and relations between $HH_*(\mathcal{C})$ and $HH^*(\mathcal{C})$ should be reflected in a relation between $\mathcal{DA}(\mathcal{C})$ and $\mathcal{D}\Delta(\mathcal{C})$. We will proceed in this direction elsewhere. At present, the best we can do is to make the following hopeful observation:

- the category $\text{Sec}_{\text{cart}}(\mathcal{C}^{\#})$ is naturally a *braided* tensor category over k .

The reason for this is very simple: if one writes out explicitly the definition of $\text{Sec}_{\text{cart}}(\mathcal{C}^{\#})$ along the lines of Lemma 2.2, one finds that it coincides on the nose with the Drinfeld double of the tensor category \mathcal{C} .

References

- [BD] I. BUCUR, A. DELEANU, *Introduction to the theory of categories and functors*, Interscience Publication John Wiley & Sons, Ltd., London-New York-Sydney 1968.
- [C1] A. CONNES, *Non-commutative differential geometry, I, II*, Publ. IHES, **62** (1985), 257–360.
- [C2] A. CONNES, *Cohomologie cyclique et foncteur Ext^n* , Comptes Rendues Ac. Sci. Paris Sér. A-B, **296** (1983), 953–958.
- [FT] B. FEIGIN, B. TSYGAN, *Additive K-Theory*, in Lecture Notes in Math. **1289** (1987), 97–209.
- [GM1] S. GELFAND, YU. MANIN, *Methods of homological algebra*, Nauka Publishers, Moscow, 1988 (in Russian).
- [GM2] S. GELFAND, YU. MANIN, *Homological algebra*, Itogi vol. **38** (Algebra V), VINITI, Moscow, 1989 (in Russian).
- [Ge] E. GETZLER, *Cartan homotopy formulas and the Gauss-Manin connection in cyclic homology*, in *Quantum deformations of algebras and their representations (Ramat-Gan, 1991/1992; Rehovot, 1991/1992)*, Israel Math. Conf. Proc. **7**, Bar-Ilan Univ., Ramat Gan, 1993, 65–78.
- [Go] T. GOODWILLIE, *Cyclic homology, derivations, and the free loop space*, Topology **24** (1985), 187–215.
- [Gr] A. GROTHENDIECK, *Exposé VI: Catégories fibré et descente*, in *SGAI: Revêtements étales et groupe fondamental*, Lecture Notes in Math., **224**, Springer, Berlin; 145–194.
- [Ka] D. KALEDIN, *Non-commutative Hodge-to-de Rham degeneration via the method of Deligne-Illusie*, Pure. Appl. Math. Q., **4** (2008), no. 3, part 2, 785–875.
- [L] J.-L. LODAY, *Cyclic Homology*, second ed., Springer, 1998.

- [TT] D. TAMARKIN, B. TSYGAN, *The ring of differential operators on forms in noncommutative calculus*, in *Graphs and patterns in mathematics and theoretical physics*. Proc. Sympos. Pure Math. **73**, AMS, Providence, RI, 2005; 105–131.
- [Ts] B. TSYGAN, *Homology of Lie algebras over rings and Hochschild homology*, *Uspekhi Mat. Nauk*, **38** (1983), 217–218.
- [W] C. WEIBEL, *Cyclic homology for schemes*, Proc. AMS **124** (1996), 1655–1662.



<http://www.springer.com/978-0-8176-4746-9>

Algebra, Arithmetic, and Geometry
Volume II: In Honor of Yu. I. Manin
Tschinkel, Y.; Zarhin, Y. (Eds.)
2009, XII, 704 p. 15 illus., Hardcover
ISBN: 978-0-8176-4746-9
A product of Birkhäuser Basel