

Piotr K. Janicki

## Key Points

- Individual pain variability and differences in the efficacy of analgesic drugs are genetically controlled.
- Drug-metabolizing enzymes represent a major target of current effort to identify associations between individuals' analgesic drug response and genetic profile.
- Genetic variants in other candidate genes influencing drug effector sites, such as those encoding receptors, transporters, and other molecules important for pain transmission represent another, less well-defined target.
- The pharmacogenomics-based approach to pain management represents a potential tool to improve the effectiveness and the side effect profile of therapy; however, well-designed prospective studies are needed to demonstrate superiority to conventional dosing regimes.

The goal of pharmacogenomics is to use information provided by advances in human genetics to identify patients at risk for significantly altered response during pharmacotherapy. The field of pharmacogenomics represents the major drive behind the introduction of the concept of *personalized medicine* in which the medical treatment is customized according to the individual patient genomic signature [1].

## Background

Association of genome variability with increased or decreased pain, or modified effects of analgesics, has demonstrated that pain therapy is subject to pharmacogenomics [2–6].

There are two major components of pain management and pharmacogenomics (see Fig. 2.1). The use of genetic information from basic science and clinical studies to examine the impact of genetic variability on factors modulating the risk of developing pain, its clinical course, and intensity is called *functional pain genomics*. Functional pain genomics aims to discover the biologic function of particular genes and to uncover how a set of genes and their products work together in regulating the response to pain.

The second, more traditional, and better established component of pain related genomics is called *pharmacogenomics of pain management* and aims to characterize how genetic variations contribute to an individual's sensitivity and response to a variety of drugs important to pain management practice. Pharmacogenomics is traditionally divided into two parts describing genetic variants influencing pharmacokinetics and pharmacodynamics.

The molecular basis for the observed variability in patient response is defined by different forms of the detected genetic variants. These variants, consisting of the interindividual differences in the DNA sequences, produce the individual *phenotypes* of the human being. There are many different types of genetic variants (see Fig. 2.2). The most common (more than ten million types known so far) are single *nucleotide polymorphisms (or SNP)*, which represent a point mutation


## Introduction

Medicine has been continuously challenged, as well as stimulated, by the extraordinary variability in patient response to pharmacotherapy. The new age of identification of risk factors associated with pharmacotherapy using the methods of molecular medicine focuses on generating predictions regarding clinical outcome on the basis of each individual's unique DNA sequence. This new field has been coined *pharmacogenomics*.

P.K. Janicki, M.D., Ph.D., DSci, DABA  
Department of Anesthesiology,  
Pennsylvania State Milton S. Hershey Medical Center,  
500 University Drive, Mailcode H187, Hershey, PA 17033, USA

Laboratory of Perioperative Genomics, Pennsylvania State University  
College of Medicine, Hershey, PA, USA  
e-mail: pjanicki@hmc.psu.edu

**Fig. 2.1** Framework of genetic background influencing the response to analgesic drugs



Relative frequency	Genetic variants	Number of base pairs (bp) affected
	Single nucleotide polymorphism = SNP aatgc...>aatac	1 (single) bp
	Deletion/insertion atgc...>atgac	1–1000s bp
	Variable number of tandem repeats = VNTR atg<u>cgcg</u>cg...>atg<u>cgcg</u>cgcgcgcg	Repeats of 2 bp
	Microsatellites atgcaacc...>atgcaaccgcaacc	Repeats of 6–8 bp
	Minisatellites	Repeats of 20–30 bp
	Gene copy numbers — — —>— — —	Repeats of whole genes
	Chromosomal aberrations	Repeats of chromosomes

**Fig. 2.2** Types of genetic variants taking part in modifying pain phenotype

(change of one base) in the DNA fragments. Other allelic mutations include insertion or deletion of a single base (*indels*), multiple, continuous repeats of 2–4 bases (*variable number of tandem repeats or VNTR*); repeats of longer DNA fragments (*micro- and mini-satellites*); *copy number vari-*

*ants* (CNV, deletion or multiplication of large, >1,000 bases fragments of chromosomes); and finally *chromosomal aberrations*. The genetic variants may produce alterations in the protein's function through either changes in the protein expression or its structure.

## Functional Genomics of Pain

Pain as a complex trait is expected to have a polygenic nature shaped by the environmental pressures. Identification of specific genetic elements of pain perception promises to be one of the key elements for creating novel and individualized pain treatments. It was demonstrated previously that both rare deleterious genetic variants and common genetic polymorphisms are mediators of human pain perception and clinical pain phenotypes [7, 8]. A higher or lower intensity of pain is very likely to require higher or lower doses of analgesics for efficient pain management. The genetic control of human pain perception and processing is therefore likely to modulate analgesic therapy.

The complete inability to sense pain in an otherwise healthy individual is a very rare phenotype. At present, five types of congenital insensitivity to pain (or HSAN=hereditary sensory and autonomic neuropathy) were identified which are caused by mutations in five different genes [9].

Recently, several new genomic mutations were identified which are described as “channelopathy-associated insensitivity to pain” [10] which are characterized by complete and selective inability to perceive any form of pain. It includes mutations in the alpha-subunit of sodium channel  $Na_v1.7$  (SCN9A), causing the loss of function in this specific form of sodium channel [10, 11]. By contrast, mutations in SCN9A that leads to excessive channel activity trigger activation of pain signaling in humans and produce primary erythralgia (more frequently used term is erythromelalgia), which is characterized by burning pain in response to exposure to mild warmth [12, 13]. Mutations in this gene also produce a rare condition referred to as “paroxysmal extreme pain disorder,” which is characterized by rectal, ocular, and submandibular pain [14].

These syndromes probably have no importance in the everyday clinical pain management as they are very rare, and the affected people probably do not require pain therapy (with exception of erythromelalgia which causes severe pain that is considered a true pain-related emergency). However, defining the molecular causes for hereditary insensitivity to pain may serve as an important source of information to find new targets for analgesic drugs. This assumption was confirmed in the recently published study, in which the authors after investigating 27 common polymorphisms in the SCN9A gene found out that the minor A allele of the SNP rs6746030 was associated with an altered pain threshold and the effect was mediated through C-fiber activation [15]. They concluded that individuals experience differing amounts of pain, per nociceptive stimulus, on the basis of their SCN9A rs6746030 genotype.

Pain in the average population is controlled by fairly frequent genetic variants (allelic frequency > 10 %). Each of them, however, modifies the pain phenotype to only modest


degree, and in the majority of cases, the evidence for their involvement in the efficacy of analgesics is either lacking or remains controversial [7, 8]. The involvement of common variants of the opioid receptors, kappa and mu, are discussed below in the part describing pharmacodynamic modifications of activity of opioid analgesics. A variant of third type of opioid receptor, delta, has been associated with lower thermal pain intensity with no association, so far, with the efficacy of opioid analgesics [16].

GTP cyclohydrolase (GCH1), recently implicated in shaping pain responses in humans, regulates production of tetrahydrobiopterin (BH4), an essential factor for the synthesis of dopamine, serotonin, and nitric acid. Tegeder et al. discovered a haplotype associated with reduction of experimental pain in normal volunteers and a favorable outcome with regard to long-term pain reduction that underwent pain (did you mean “a painful surgery”?) surgery [17]. In another study, Tegeder et al. showed that carriers of the particular GCH1 haplotype had higher pain threshold to mechanical and thermal pain following capsaicin sensitization [18]. However, Kim and Dionn and Lazarev et al. failed to replicate significant associations between the same GCH1 genomic variants and pain responses, both in assessment of experimental pain and postoperative pain after dental surgery, as well chronic pancreatic pain [19, 20]. Conversely, the most recent study confirmed again that the five previously identified GCH1 SNPs were profoundly affecting the ratings of pain induced by capsaicin in healthy human volunteers [21]. It was also suggested that the carriers of this particular GCH1 haplotype (which may be responsible for the decreased function of GCH1) display delayed need for pain therapy [2, 22].

## Pharmacogenomics of Pain Therapy and Its Usefulness in Clinical Practice

Pharmacogenomics of pain management represents the most familiar area of practical pain genomics. It includes several examples of genomic variations, dramatically changing response to analgesic drugs through either change in their metabolism or receptor targets.

The current list of genetic polymorphisms which may affect the action of analgesic drugs is quite long and appears to be growing rapidly. The best known mechanisms involved in the altered effects of analgesics involve polymorphic changes in its metabolism. In this respect, three major mechanisms have been identified, involving genetic variations in the metabolic activation of the analgesics administered as an inactive or less active prodrug, variations in the metabolic degradation of the active components, and variations in its transmembrane transport.



**Fig. 2.3** Opioid analgesics influenced by polymorphic CYP2D6 metabolism (3-O-demethylation)

### Genetic Variations in the Prodrug Activation

The better known example involves polymorphisms in genes of the liver isoforms of the cytochrome P450 system (CYP) [23]. In particular, the most well-characterized *CYP2D6* polymorphism is responsible for the considerable variation in the metabolism (and clinical responses) of drugs from many therapeutic areas, including several analgesics (Fig. 2.3) [24–26]. More than 100 *CYP2D6* alleles have been identified, ranging from nonsynonymous mutations to SNPs that either alter RNA splicing or produce deletions of the entire gene [27]. Of these, \*3, \*4, and \*8 are nonfunctional, \*9, \*10, and \*41 have reduced function, and \*1, \*2, \*35, and \*41 can be duplicated, resulting in greatly increased expression of functional *CYP2D6*. There are also interethnic differences in the frequencies of these variant alleles. Allele combinations determine phenotype: two nonfunctional = poor metabolizer (PM); at least one reduced functional = intermediate metabolizer (IM); at least one functional = extensive metabolizers (EM); and multiple copies of a functional and/or allele with promoter mutation = ultrarapid metabolizer (UM). The most recent update of *CYP2D6* nomenclature and terminology could be found on home Web page of the Human Cytochrome P450 (CYP) Allele Nomenclature at <http://www.cypalleles.ki.se/>.

Codeine and other weak opioids are extensively metabolized by polymorphic *CYP2D6* which regulates its O-demethylation to more potent metabolites (e.g., after a single oral dose of 30 mg codeine, 6% is eventually transformed

to morphine). The clinical analgesic effect of codeine is mainly attributed to its conversion to morphine, which has a 200 times higher affinity and 50 times higher intrinsic activity at MOR than codeine itself [28]. Since *CYP2D6* is genetically highly polymorphic, the effects of codeine are under pharmacogenetic control.

Genetically, altered effects of codeine may occur in subjects with either decreased, absent, or highly increased *CYP2D6* activity when compared with the population average [29, 30]. Decreased or absent *CYP2D6* activity in PMs causes production of only very low or absent amount of morphine after codeine administration. The ultrafast metabolizers (UM) produce on the other hand excessive amount of morphine after typical dose of codeine. Roughly, one out of seven Caucasians is at risk of either failure or toxicity of codeine therapy due to extremely low or high morphine formation, respectively. Recent case reports of codeine fatalities highlighted that the use of this weak opioid, particularly in young children, is associated with a substantial risk in those subjects displaying UM genotype [31–37]. The polymorphic variants in the *CYP2D6* system are responsible for some but not all variability observed after codeine administration. The other causes for the observed high variability in codeine efficacy include both polymorphisms in other genes involved in opioid expression or trafficking, as well as nongenetic factors. In addition to differences in codeine metabolism between EM and PM, the differences between EM of various ethnicities have also been highlighted. The Chinese EM reported having a lower rate of codeine O-demethylation when compared with the Caucasian EM, because of the much higher frequency (50%) of the \*10 (reduced function) allele in the Chinese [27].

Other popular analgesic drug which depends on activation by the *CYP2D6* includes *tramadol*. Tramadol is a mu-opioid receptor (MOR) agonist but has a lower affinity at MORs than its active metabolite O-desmethyltramadol. Tramadol itself has weak analgesic activity which becomes evident when *CYP2D6* is blocked and also acts through non-opioid-dependent mechanisms which involve serotonin- and noradrenaline-mediated pain inhibition originating in brain stem. The analgesic activity of tramadol is strongly modulated by *CYP2D6* activity. The analgesic activity on experimental pain is reduced in *CYP2D6* PMs, a finding later confirmed in pain patients [38–42]. It is interesting to note that the pharmacokinetics of tramadol (which is administered as racemic substance containing equal amount of (–) and (+) optical isomers producing analgesia by a synergistic action of its two enantiomers and their metabolites) is enantioselective in *CYP2D6* poor and extensive metabolizers, meaning that the production of either optical isomer differs depending on the metabolic status [41, 43, 44]. The clinical significance of this finding for pain management remains to be further explored.

As far as other CYP2D6 substrates with active analgesic metabolites are concerned (see Fig. 2.3), the evidence for variation in its analgesic with altered CYP2D6 function is less evident when compared with codeine or tramadol. These examples are either negative, such as for dihydrocodeine [45, 46], or explained at a nongenetic level, such as for oxycodone. In other cases, the evidence is based only on animal studies, such as hydrocodone, or restricted to single case reports for inadequate activity of oxycodone [3].

*Tilidine* (an opioid analgesic) is activated to active metabolite nortilidine, and *parecoxib* (an NSAID) is activated into valdecoxib by CYP3A system. This enzyme is phenotypically highly variable, but only a minor part of this variability can be attributed to genetics [2]. Individuals with at least one CYP3A5\*1 allele copy produce fully active copy of CYP3A5 enzyme; however, the majority of Caucasians have no active CYP3A5 due to a premature stop codon.

### Genetic Variations in the Elimination of Analgesic Drugs

Many of the opioids contain hydroxyl group at position 6, and the potent opioids have a hydroxyl at position 3 of the 4,5-methoxymorphinan structure. The glucuronidation of morphine, codeine, buprenorphine, dihydrocodeine, dihydromorphine, hydromorphone, dihydromorphine, oxymorphone, as well as opioid receptor antagonists (naloxone and naltrexone) is mainly mediated by the uridine diphosphate (UDP) glucuronyltransferase (UGT)2B7 [3]. Similar to CYP genes, the UGT2B7 gene is also polymorphic, although less than 20 allelic variants have been identified. The main proportion of morphine is metabolized to morphine-6-glucuronide, M6G (approximately 70 %), and to lesser degree to morphine-3-glucuronide (M3G). Both metabolites are active, with effects opposite to each other, consisting in excitation and anti-analgesia for M3G and in typical opioid agonist effects for M6G. Despite the role of UGT2B7 in the formation of M6G and M3G, the clinical effect of the UGT2B7\*2 (268Y) variant has only produced conflicting results so far. Different variants in the 5' untranslated region of UGT2B7 are associated with reduced M6G/morphine ratios in patients. In addition, it was reported that UGT2B7\*2/\*2 genotypes and CYP2D6 UM phenotypes were associated with severe neonatal toxicity after breast-feeding and oral ingestion of opioids. The above preliminary data indicate that the consequences of UGT variants were so far restricted to alterations of plasma concentrations, while none of the UGT variants alone have been associated with the altered efficacy of opioid analgesics [3, 4].

The increased enzyme activity associated with the CYP3A5\*1 allele may cause accelerated elimination of CYP3A substrates, such as alfentanil, fentanyl, or sufentanil.

However, positive associations of CYP3A polymorphisms with analgesic actions have not been reported so far. The CYP3A5 genotype did not affect the systemic or apparent oral clearance as well as the pharmacodynamics of alfentanil and levomethadone [47, 48].

In addition to CYP2D6 and CYP2A5, there is also clinical evidence about the involvement of other CYP systems in the metabolism of frequently used nonsteroid anti-inflammatory drugs (NSAIDs). Human CYP2C9 metabolizes numerous drugs (e.g., warfarin, oral sulfonylurea hypoglycemics, antiepileptics, and others) [49]. In addition, CYP2C9 polymorphism might play a significant role in the analgesic efficacy and toxicity of traditional NSAIDs, for example, diclofenac, ibuprofen, naproxen, tenoxicam, and piroxicam, as well as selective COX-2 inhibitors such as celecoxib and valdecoxib [50]. More than 33 variants and a series of subvariants have been identified for CYP2C9 to date. The two missense mutations, CYP2C9\*2 (rs1799853) and CYP2C9\*3 (rs1057910), yield enzymes with decreased activity [51]. These alleles are mainly present in Caucasians, while their frequency is lower in African and Asian subjects. More than twofold reduced clearance after oral intake of celecoxib was observed in homozygous carriers of CYP2C9\*3 compared with carriers of the wild-type genotype CYP2C9\*1/\*1 [52]. Similarly, ibuprofen-mediated inhibition of COX-1 and COX-2 is significantly decreased (by 50 %) in carriers of two CYP2C9\*3 alleles [53]. Further investigations demonstrating the relevance of the CYP2C9\*3 allele for naproxen, tenoxicam, piroxicam, and lornoxicam pharmacokinetics have been also published [4, 54]. Although CYP2C9 is the major determinant of clearance, it is necessary to also consider CYP2C8 genotype, as it contributes to some smaller extent in NSAIDs metabolism. In the study performed in healthy volunteers, it was demonstrated that metabolism of diclofenac was significantly slower in individuals carrying CYP2C8\*3 (rs10509681) or CYP2C8\*4 (rs1058930) allele than in those homozygous for the wild-type allele [55].

Whereas numerous clinical trials have demonstrated the impact of CYP2C9\*3 on therapy with Coumadin, less information is available on the CYP2C9 genotype-related efficacy of NSAIDs in the pain management. Some publications focus, however, on the incidence and severity of adverse effects (e.g., gastrointestinal (GI) bleeding, effects on coagulation). It was that the combined presence of CYP2C8\*3 and CYP2C9\*2 was a relevant determinant in the risk of developing GI bleeding in patients receiving NSAIDs metabolized by CYP2C8/9 [56]. Similar results were also presented by Agundez et al. [57]. However, to date, the study results from other authors are conflicting, with several other trials reporting no association [58, 59]. More studies are clearly necessary to confirm the relevance of CYP2C8/9 genotype with increased incidence of GI bleeding.

Another typical adverse effect is the influence of classical NSAIDs on coagulation. The risk of altered coagulation was substantially increased in patients with either CYP2C9\*3 and CYP2C9\*3 (mentioned twice) genotypes taking Coumadin together with NSAIDs which are known CYP2C9 substrates [60].

### Genetic Variations in the Transmembrane Transport of Analgesics

P-glycoprotein (P-gp) coded by the ATP-binding cassette subfamily (ABCB1)/multidrug resistance (MDR1) gene is mainly located in organs with excretory functions (e.g., liver, kidneys). It is also expressed at the blood-brain barrier where it forms an outward transporter. Therefore, functional impairment of P-gp-mediated drug transport may be expected to result in increased bioavailability of orally administered drugs, reduced renal clearance, or an increased brain concentration of its substrates. Some opioids are P-gp substrates. The ABCB1 3435 C>T variant (rs1045642) is associated with decreased dosage requirements in opioids that are P-gp substrates, as assessed in outpatients. Moreover, a diplotype consisting of three polymorphic positions in the ABCB1 gene (1236TT-rs1128503, 2677TT-rs2032582, and 3435TT) is associated with increased susceptibility to respiratory depression caused by fentanyl in Korean patients [61]. The results suggest that analysis of ABCB1 polymorphisms may have clinical relevance in the prevention of respiratory suppression by intravenous fentanyl or to anticipate its clinical effects. With the OPRM1 118 A>G variant (see below), the ABCB1 3435 C>T predicted the response to morphine in cancer patients with a sensitivity close to 100 % and a specificity of more than 70 % [62]. Trials in patients suffering from chronic and cancer pain had shown decreased opioid consumption in carriers of the 3435T allele [63, 64]. Finally, methadone analgesia may be subject to P-gp pharmacogenetic modulation. The pupillary effects of orally administered methadone are increased following the pharmacological blockade of P-gp by quinidine, and the methadone dosing for heroin substitution can be decreased in carriers of ABCB1 variants associated with decreased transporter expression, for example, ABCB1 2435 C>T and others [65, 66].

### Pharmacodynamics of Pain Therapy

The alterations in effects of analgesics may also result from pharmacodynamic interferences, consisting of altered receptor binding, activation or signaling mechanisms, or of altered expression of the drug's target, such as opioid receptors or cyclooxygenases. Genetic factors have been found to act via any of these mechanisms.

### Opioid Receptors

The mu-opioid receptor (MOR) is part of the family of several types of opioid receptors which are 7-transmembrane domain, G-protein-coupled receptors (GPCR), and inhibit cellular activity. MOR is clinically most relevant target of opioid analgesics. The OPRM1 gene coding for MOR in humans is highly polymorphic, with excess of 1,800 SNPs listed in the current edition (2010) of the NCBI SNP database (<http://www.ncbi.nlm.nih.gov/snp>). Coding mutations affecting the third intracellular loop of MOR (e.g., 779 G>A, 794 G>A, 802 T>C) result in reduced G-protein coupling, receptor signaling, and desensitization, leading to an expectation that opioids should be almost ineffective in patients carrying those polymorphisms. However, these polymorphisms are extremely rare (<0.1 % population) and are therefore restricted to very rare single cases.

Evidence for a function of OPRM1 variants with allelic frequencies >5 % is sparse, except for the 118 A>G polymorphism (rs1799971). This SNP causes an amino acid exchange of the aspartate with an asparagine at position 40 of extracellular part of MOR, deleting one of a putative glycosylation sites. This change can cause altered expression of MOR or its signaling [67–69]. The OPRM1 118 A>G polymorphism has an allele frequency of 8–17 % in Caucasians and considerably higher in Asians, with a frequency of 47 % reported from Japan. It is also worth noting that the frequency of homozygotes for the GG allele is by much higher in Asian population with only very rare (<1 %) occurrence in Caucasian population [70]. The data obtained so far with the OPRM1 118 A>G polymorphism have been controversial [71]. The molecular changes associated with SNP 118 A>G translate to a variety of clinical effects (predominantly decrease) of many opioids in experimental settings and clinical studies [72–81]. The consequences of the SNP 118 A>G have consistently been related to a decrease in opioid potency for pupil constriction (e.g., for morphine, M6G, methadone). For analgesia, the SNP decreases the concentration-dependent effects of alfentanil on experimental pain. Specifically, the variant decreases the effect of opioids on pain-related activation mainly in those regions of the brain that are processing the sensory dimension of pain including the primary and secondary somatosensory cortex and posterior insular cortex [82]. In clinical settings, greater postoperative requirements of alfentanil and morphine have been reported for carriers of the variant, and higher concentrations of alfentanil of M6G were needed to produce analgesia in experimental pain models [2, 48, 83–87]. It should be noted that other studies described only moderate to no significant effects of the OPRM1 118 A>G polymorphism on opioid requirements or pain relief. Several studies did not demonstrate any association between OPRM1 variant and analgesic needs [88–92]. Contradictory results were

reported by Landau et al. who investigated the influence of OPRM1 118 A>G polymorphism on the analgesic effectiveness of fentanyl in females after its intrathecal administration during labor and delivery. The analgesic requirements in this study were increased in homozygous carriers of AA allele, the opposite effect compared with most other studies [93]. In the chronic pain patients, it was reported that in the high-quartile opioid utilization group, the homozygous carriers of the minor allele required significantly higher opioid doses than the carriers of the minor allele [91]. In another studies, GG homozygote patients were characterized by higher morphine consumption than carriers of the major AA allele [76, 94]. In summary, an influence of OPRM1 genetic variants on opioid requirements and degree of pain relief under opioid medication has been demonstrated in some studies; however, this could not be replicated in all subsequent investigations. Patients stratification; a low number of patients with the GG genotype (in particular in studies performed in Caucasian populations); presence of multiple, uncontrolled co-variables influencing the phenotype; and a clinically questionable reduction in opioid consumption are some major concerns. The requirements of high opioid doses may in part reflect an addiction component or a higher/faster rate of tolerance development in certain pain patients. It was reported that OPRM1 A118G polymorphism is a major determinant of striatal dopamine responses to alcohol. Social drinkers recruited based on OPRM1 genotype were challenged in separate sessions with alcohol and placebo under pharmacokinetically controlled conditions and examined for striatal dopamine release using positron emission tomography and [(11)C]-raclopride displacement. A striatal dopamine response to alcohol was restricted to carriers of the minor 118G allele. Based on the results of this study, it was concluded that OPRM1 A118G variation is a genetic determinant of dopamine responses to alcohol, a mechanism by which it likely modulates alcohol reward [95].

In addition, the most recent study seems to suggest that some of the effect of SNP A>G could be explained by the linkage disequilibrium with other functional SNPs located in the OPRM1 region [96]. For example, SNP rs563649 is located within a structurally conserved internal ribosome entry site in the 5'-UTR of a novel exon 13-containing OPRM1 isoforms (MOR-1K) and affects both mRNA levels and translation efficiency of these variants. Furthermore, rs563649 exhibits very strong linkage disequilibrium throughout the entire OPRM1 gene locus and thus affects the functional contribution of the corresponding haplotype that includes other functional OPRM1 SNPs. These results might provide evidence for an essential role for MOR-1K isoforms in nociceptive signaling and suggest that genetic variations in alternative OPRM1 isoforms may contribute to individual differences in opiate responses.

### **Catechol-O-Methyltransferase (COMT)**

COMT degrades catecholamine neurotransmitters such as norepinephrine, epinephrine, and dopamine. Increased dopamine concentrations suppress the production of endogenous opioid peptides. Opioid receptor expression is in turn upregulated, which has been observed with the Val158Met variant of COMT, coded by the COMT 772 G>A (rs4680) SNP in human postmortem brain tissue and in vivo by assessing radiolabeled 11C-carfentanil MOR binding [97, 98]. This variant leads to a low-function COMT enzyme that fails to degrade dopamine, which may cause a depletion of enkephalin. Patients with cancer carrying the Val158Met variant needed less morphine for pain relief than patients not carrying this variant. Finally, the variant exerts its opioid enforcing effects also in cross relation with the OPRM1 118 A>G variant [94, 97–101]. During the past decade, several new polymorphisms were identified in the COMT gene which contains at least five functional polymorphisms that impact its biological activity and associated phenotypes (including pain). The potentially complex interactions of functional variations in COMT imply that the overall functional state of the gene might not be easily deduced from genotype information alone, which presumably explains the inconsistency in the results from association studies that focus on the Val158Met polymorphism [102, 103].

### **Melanocortin 1 Receptor (MC1R)**

Nonfunctional variants of the MC1R which produces bright red hair and fair skin phenotype were associated with an increased analgesic response to kappa opioid receptors (KOR)-mediated opioid analgesia. Red-headed women required less of the KOR agonist drug – pentazocine – to reach a specific level of analgesia compared with all other groups [104, 105]. This study presented the first strong evidence for a gene-by-sex interaction in the area of pain genetics, because the authors also showed that red-headed men did not experience enhanced KOR analgesia.

### **Cyclooxygenases (COX)**

Polymorphisms in the prostaglandin endoperoxidase synthase 2 gene (PTGS2) coding for COX-2 may modulate the development of inflammation and its response to treatment with inhibitors of COXs, especially those specific for COX-2 [106]. This has been proposed for the PTGS2-765 G>C SNP (rs20417), which was reported to be associated with more than a twofold decrease in COX-2 expression [107]. By altering a putative Sp1 binding site in the promoter region of PTGS2, this gene variant was found to decrease the promoter activity by 30 % [108]. However, the controversial results were reported so far in clinical studies with this polymorphisms and different COX-2 inhibitors. The inhibitory effect of celecoxib on COX-2 was not associated with the presence of this variant in volunteers [109];

conversely, significantly decreased analgesic effects of rofecoxib were observed in the homozygous carriers of this variant [110].

### Future Direction of Pharmacogenomics in Pain Treatment

The influence of different genetic variants on analgesic requirements and degree of pain relief has been demonstrated in some studies; however, there is relatively less information available about the interactions between these variants. Each of the genetic variants investigated up to now seems to contribute in a modest way to the modulation of analgesic response [111]. However, a global approach investigating multiple possible variables within one trial has not been performed. After more than a decade of identifying genetic associations, the current challenge is to intensify compilation of this information for precisely defined clinical settings for which improved pain treatment is possible.

The current knowledge about the impact of genetics in the pain management is based on the association studies. In contrast to traditional family or pedigree-based studies (linkage analysis), in this type of studies, two cohorts of unrelated patients (with and without the observed phenotype, i.e., changes in the efficacy of analgesics) are compared in respect to the frequency of different genetic variants (adjusted for other known risk factors and for environmental differences). Candidate-gene association studies are focused on selected genes which are thought to be relevant for a specific observed outcome.

The alternative to targeted association studies are genome-wide association studies (GWAS). In this type of studies, there is no a priori hypothesis about the gene candidates. Instead, the microarray-based genomic scans are performed throughout the whole genome in order to find all SNPs possibly associated with observed phenotypic changes in the cohorts of patients with investigated traits (and controls). The modern microarray platforms allow for the cost-effective, parallel analysis of approximately one million genomic variants in one sample (or pooled samples) and, using sophisticated computer strategy, enable finding the most relevant statistical associations between control and affected patients. The main advantage of GWAS is that it is an unbiased hypothesis-free approach. In contrast to other areas of medicine, the GWAS approach lags behind in pain genomics, but the next few years should bring about the results of several studies currently being performed in the area of pain medicine. One of the first pain pharmacogenomic studies using GWAS technology was recently published by Kim et al. and demonstrated association of minor allele variant in a zinc finger protein (ZNF429) gene with delayed onset of action of ketorolac in the oral surgery patients [112].

**Table 2.1** List of the most common analgesic drugs and polymorphic genes for which some evidence exists that the pharmacokinetics and/or pharmacodynamics of these analgesic drugs are modulated by functional genetic variants

Analgesic drug	Genes
<i>Opioid analgesics</i>	
Codeine	CYP2D6, UGT2B7, ABCB1, OPRM1
Pentazocine	MC1R
Tramadol	CYP2D6
Morphine	UGT2B7, ABCB1, COMT, OPRM1, CGH1
Methadone	CYP2D6, UGT2B7, ABCB1, OPRM1
Tilidine	CYP3A
Dihydrocodeine, hydrocodone, oxycodone	CYP2D6, ABCB1, COMT, OPRM1
<i>NSAIDs</i>	
Ibuprofen	CYP2C9
Diclofenac	CYP2C9
Naproxen	CYP2C9
Valdecoxib	CYP2C9, PTGS2
Celecoxib	CYP2C9, PTGS2
Parecoxib	CYP3A, CYP2C9, PTGS2

### Summary

In summary, genetics continues to make rapid progress in terms of technology and understanding, but there are still, as yet, no large randomized, multicenter controlled trials to support the use of widespread genetic screening to predict an individual's response to pain medication (Table 2.1) [113]. Despite intensive research, genetics-based personalized pain therapy has yet to emerge. Monogenetic heredity of pain conditions seems to be restricted to very rare and extreme phenotypes, whereas common phenotypes are very complex and multigenetic. Many common variants, of which only a fraction have been identified so far, produce only minor effects that are sometimes partly canceled out. For most clinical settings and analgesic drug effects, common genetic variants cannot yet be used to provide a relevant prediction of individual pain and analgesic responses. However, genetics has some potential practical uses: CYP2D6, MC1R, and potentially PTGS2 could provide guidance on the right choice of analgesics. We still have a way to go before genetic screening becomes a routine practice and much further still before the contribution of gene-environment interactions is fully realized. However, continued identification of genotypes which are predictive of efficacy of pain management may not only further our understanding of the pain mechanisms but also potentially help discover new potential molecular targets for pain therapy.

## References

- Eichelbaum M, Ingelman-Sundberg M, Evans WE. Pharmacogenomics and individualized drug therapy. *Annu Rev Med.* 2006;57:119–37.
- Lotsch J, Geisslinger G, Tegeder I. Genetic modulation of the pharmacological treatment of pain. *Pharmacol Ther.* 2009;124:168–84.
- Somogyi AA, Barratt DT, Collier JK. Pharmacogenetics of opioids. *Clin Pharmacol Ther.* 2007;81:429–44.
- Stamer UM, Zhang L, Stuber F. Personalized therapy in pain management: where do we stand? *Pharmacogenomics.* 2010;11:843–64.
- Lacroix-Fralish ML, Mogil JS. Progress in genetic studies of pain and analgesia. *Annu Rev Pharmacol Toxicol.* 2009;49:97–121.
- Landau R. One size does not fit all: genetic variability of mu-opioid receptor and postoperative morphine consumption. *Anesthesiology.* 2006;105:235–7.
- Diatchenko L, Nackley AG, Tchivileva IE, Shabalina SA, Maixner W. Genetic architecture of human pain perception. *Trends Genet.* 2007;23:605–13.
- Fillingim RB, Wallace MR, Herbstman DM, Ribeiro-Dasilva M, Staud R. Genetic contributions to pain: a review of findings in humans. *Oral Dis.* 2008;14:673–82.
- Nagasako EM, Oaklander AL, Dworkin RH. Congenital insensitivity to pain: an update. *Pain.* 2003;101:213–9.
- Cox JJ, Reimann F, Nicholas AK, et al. An SCN9A channelopathy causes congenital inability to experience pain. *Nature.* 2006;444:894–8.
- Goldberg YP, MacFarlane J, MacDonald ML, et al. Loss-of-function mutations in the Nav1.7 gene underlie congenital indifference to pain in multiple human populations. *Clin Genet.* 2007;71:311–9.
- Waxman SG. Neurobiology: a channel sets the gain on pain. *Nature.* 2006;444:831–2.
- Waxman SG, Dib-Hajj SD. Erythromelalgia: a hereditary pain syndrome enters the molecular era. *Ann Neurol.* 2005;57:785–8.
- Fertleman CR, Baker MD, Parker KA, et al. SCN9A mutations in paroxysmal extreme pain disorder: allelic variants underlie distinct channel defects and phenotypes. *Neuron.* 2006;52:767–74.
- Reimann F, Cox JJ, Belfer I, et al. Pain perception is altered by a nucleotide polymorphism in SCN9A. *Proc Natl Acad Sci USA.* 2010;107:5148–53.
- Kim H, Mittal DP, Iadarola MJ, Dionne RA. Genetic predictors for acute experimental cold and heat pain sensitivity in humans. *J Med Genet.* 2006;43:e40.
- Tegeder I, Costigan M, Griffin RS, et al. GTP cyclohydrolase and tetrahydrobiopterin regulate pain sensitivity and persistence. *Nat Med.* 2006;12:1269–77.
- Tegeder I, Adolph J, Schmidt H, Woolf CJ, Geisslinger G, Lotsch J. Reduced hyperalgesia in homozygous carriers of a GTP cyclohydrolase 1 haplotype. *Eur J Pain.* 2008;12:1069–77.
- Kim H, Dionne RA. Lack of influence of GTP cyclohydrolase gene (GCH1) variations on pain sensitivity in humans. *Mol Pain.* 2007;3:6.
- Lazarev M, Lamb J, Barmada MM, et al. Does the pain-protective GTP cyclohydrolase haplotype significantly alter the pattern or severity of pain in humans with chronic pancreatitis? *Mol Pain.* 2008;4:58.
- Campbell CM, Edwards RR, Carmona C, et al. Polymorphisms in the GTP cyclohydrolase gene (GCH1) are associated with ratings of capsaicin pain. *Pain.* 2009;141:114–8.
- Lotsch J, Klepstad P, Doehring A, Dale O. A GTP cyclohydrolase 1 genetic variant delays cancer pain. *Pain.* 2010;148:103–6.
- Ingelman-Sundberg M, Sim SC, Gomez A, Rodriguez-Antona C. Influence of cytochrome P450 polymorphisms on drug therapies: pharmacogenetic, pharmacoepigenetic and clinical aspects. *Pharmacol Ther.* 2007;116:496–526.
- Wang B, Yang LP, Zhang XZ, Huang SQ, Bartlam M, Zhou SF. New insights into the structural characteristics and functional relevance of the human cytochrome P450 2D6 enzyme. *Drug Metab Rev.* 2009;41:573–643.
- Zhou SF. Polymorphism of human cytochrome P450 2D6 and its clinical significance: Part I. *Clin Pharmacokinet.* 2009;48:689–723.
- Zhou SF, Liu JP, Lai XS. Substrate specificity, inhibitors and regulation of human cytochrome P450 2D6 and implications in drug development. *Curr Med Chem.* 2009;16:2661–805.
- Zanger UM, Raimundo S, Eichelbaum M. Cytochrome P450 2D6: overview and update on pharmacology, genetics, biochemistry. *Naunyn Schmiedeberg Arch Pharmacol.* 2004;369:23–37.
- Mignat C, Wille U, Ziegler A. Affinity profiles of morphine, codeine, dihydrocodeine and their glucuronides at opioid receptor subtypes. *Life Sci.* 1995;56:793–9.
- Thorn CF, Klein TE, Altman RB. Codeine and morphine pathway. *Pharmacogenet Genomics.* 2009;19:556–8.
- Zhou SF. Polymorphism of human cytochrome P450 2D6 and its clinical significance: part II. *Clin Pharmacokinet.* 2009;48:761–804.
- Ciszkowski C, Madadi P, Phillips MS, Lauwers AE, Koren G. Codeine, ultrarapid-metabolism genotype, and postoperative death. *N Engl J Med.* 2009;361:827–8.
- Gasche Y, Daali Y, Fathi M, et al. Codeine intoxication associated with ultrarapid CYP2D6 metabolism. *N Engl J Med.* 2004;351:2827–31.
- Koren G, Cairns J, Chitayat D, Gaedigk A, Leeder SJ. Pharmacogenetics of morphine poisoning in a breastfed neonate of a codeine-prescribed mother. *Lancet.* 2006;368:704.
- Madadi P, Koren G. Pharmacogenetic insights into codeine analgesia: implications to pediatric codeine use. *Pharmacogenomics.* 2008;9:1267–84.
- Madadi P, Koren G, Cairns J, et al. Safety of codeine during breastfeeding: fatal morphine poisoning in the breastfed neonate of a mother prescribed codeine. *Can Fam Physician.* 2007;53:33–5.
- Madadi P, Ross CJ, Hayden MR, et al. Pharmacogenetics of neonatal opioid toxicity following maternal use of codeine during breastfeeding: a case-control study. *Clin Pharmacol Ther.* 2009;85:31–5.
- Voronov P, Przybylo HJ, Jagannathan N. Apnea in a child after oral codeine: a genetic variant – an ultra-rapid metabolizer. *Paediatr Anaesth.* 2007;17:684–7.
- Enggaard TP, Poulsen L, Arendt-Nielsen L, Broesen K, Ossig J, Sindrup SH. The analgesic effect of tramadol after intravenous injection in healthy volunteers in relation to CYP2D6. *Anesth Analg.* 2006;102:146–50.
- Poulsen L, Arendt-Nielsen L, Broesen K, Sindrup SH. The hypoalgesic effect of tramadol in relation to CYP2D6. *Clin Pharmacol Ther.* 1996;60:636–44.
- Stamer UM, Lehnen K, Hothker F, et al. Impact of CYP2D6 genotype on postoperative tramadol analgesia. *Pain.* 2003;105:231–8.
- Stamer UM, Musshoff F, Kobilya M, Madea B, Hoeft A, Stuber F. Concentrations of tramadol and O-desmethyltramadol enantiomers in different CYP2D6 genotypes. *Clin Pharmacol Ther.* 2007;82:41–7.
- Stamer UM, Stuber F, Muders T, Musshoff F. Respiratory depression with tramadol in a patient with renal impairment and CYP2D6 gene duplication. *Anesth Analg.* 2008;107:926–9.
- Musshoff F, Madea B, Stuber F, Stamer UM. Enantiomeric determination of tramadol and O-desmethyltramadol by liquid chromatography-mass spectrometry and application to postoperative patients receiving tramadol. *J Anal Toxicol.* 2006;30:463–7.
- Pedersen RS, Damkier P, Broesen K. Enantioselective pharmacokinetics of tramadol in CYP2D6 extensive and poor metabolizers. *Eur J Clin Pharmacol.* 2006;62:513–21.

45. Hufschmid E, Theurillat R, Wilder-Smith CH, Thormann W. Characterization of the genetic polymorphism of dihydrocodeine O-demethylation in man via analysis of urinary dihydrocodeine and dihydromorphine by micellar electrokinetic capillary chromatography. *J Chromatogr B Biomed Appl.* 1996;678:43–51.
46. Wilder-Smith CH, Hufschmid E, Thormann W. The visceral and somatic antinociceptive effects of dihydrocodeine and its metabolite, dihydromorphine. A cross-over study with extensive and quinidine-induced poor metabolizers. *Br J Clin Pharmacol.* 1998;45:575–81.
47. Kharasch ED, Walker A, Isoherranen N, et al. Influence of CYP3A5 genotype on the pharmacokinetics and pharmacodynamics of the cytochrome P4503A probes alfentanil and midazolam. *Clin Pharmacol Ther.* 2007;82:410–26.
48. Lotsch J, Skarke C, Wieting J, et al. Modulation of the central nervous effects of levomethadone by genetic polymorphisms potentially affecting its metabolism, distribution, and drug action. *Clin Pharmacol Ther.* 2006;79:72–89.
49. Zhou SF, Zhou ZW, Huang M. Polymorphisms of human cytochrome P450 2C9 and the functional relevance. *Toxicology.* 2010;278:165–88. Epub 2009 Aug 26.
50. Rodrigues AD. Impact of CYP2C9 genotype on pharmacokinetics: are all cyclooxygenase inhibitors the same? *Drug Metab Dispos.* 2005;33:1567–75.
51. Kirchheiner J, Brockmoller J. Clinical consequences of cytochrome P450 2C9 polymorphisms. *Clin Pharmacol Ther.* 2005;77:1–16.
52. Kirchheiner J, Stormer E, Meisel C, Steinbach N, Roots I, Brockmoller J. Influence of CYP2C9 genetic polymorphisms on pharmacokinetics of celecoxib and its metabolites. *Pharmacogenetics.* 2003;13:473–80.
53. Kirchheiner J, Meineke I, Freytag G, Meisel C, Roots I, Brockmoller J. Enantiospecific effects of cytochrome P450 2C9 amino acid variants on ibuprofen pharmacokinetics and on the inhibition of cyclooxygenases 1 and 2. *Clin Pharmacol Ther.* 2002;72:62–75.
54. Bae JW, Kim JH, Choi CI, et al. Effect of CYP2C9\*3 allele on the pharmacokinetics of naproxen in Korean subjects. *Arch Pharm Res.* 2009;32:269–73.
55. Dorado P, Cavaco I, Caceres MC, Piedade R, Ribeiro V, Llerena A. Relationship between CYP2C8 genotypes and diclofenac 5-hydroxylation in healthy Spanish volunteers. *Eur J Clin Pharmacol.* 2008;64:967–70.
56. Blanco G, Martinez C, Ladero JM, et al. Interaction of CYP2C8 and CYP2C9 genotypes modifies the risk for nonsteroidal anti-inflammatory drugs-related acute gastrointestinal bleeding. *Pharmacogenet Genomics.* 2008;18:37–43.
57. Agundez JA, Garcia-Martin E, Martinez C. Genetically based impairment in CYP2C8- and CYP2C9-dependent NSAID metabolism as a risk factor for gastrointestinal bleeding: is a combination of pharmacogenomics and metabolomics required to improve personalized medicine? *Expert Opin Drug Metab Toxicol.* 2009;5:607–20.
58. Ma J, Yang XY, Qiao L, Liang LQ, Chen MH. CYP2C9 polymorphism in non-steroidal anti-inflammatory drugs-induced gastropathy. *J Dig Dis.* 2008;9:79–83.
59. Vonkeman HE, van de Laar MA, van der Palen J, Brouwers JR, Vermes I. Allele variants of the cytochrome P450 2C9 genotype in white subjects from The Netherlands with serious gastroduodenal ulcers attributable to the use of NSAIDs. *Clin Ther.* 2006;28:1670–6.
60. Visser LE, van Schaik RH, van Vliet M, et al. Allelic variants of cytochrome P450 2C9 modify the interaction between nonsteroidal anti-inflammatory drugs and coumarin anticoagulants. *Clin Pharmacol Ther.* 2005;77:479–85.
61. Park HJ, Shinn HK, Ryu SH, Lee HS, Park CS, Kang JH. Genetic polymorphisms in the ABCB1 gene and the effects of fentanyl in Koreans. *Clin Pharmacol Ther.* 2007;81:539–46.
62. Zwisler ST, Enggaard TP, Noehr-Jensen L, et al. The antinociceptive effect and adverse drug reactions of oxycodone in human experimental pain in relation to genetic variations in the OPRM1 and ABCB1 genes. *Fundam Clin Pharmacol.* 2010;24:517–24. Epub 2009 Oct 21.
63. Campa D, Gioia A, Tomei A, Poli P, Barale R. Association of ABCB1/MDR1 and OPRM1 gene polymorphisms with morphine pain relief. *Clin Pharmacol Ther.* 2008;83:559–66.
64. Lotsch J, von Hentig N, Freynhagen R, et al. Cross-sectional analysis of the influence of currently known pharmacogenetic modulators on opioid therapy in outpatient pain centers. *Pharmacogenet Genomics.* 2009;19:429–36.
65. Collier JK, Barratt DT, Dahlen K, Loennechen MH, Somogyi AA. ABCB1 genetic variability and methadone dosage requirements in opioid-dependent individuals. *Clin Pharmacol Ther.* 2006;80:682–90.
66. Levran O, O'Hara K, Peles E, et al. ABCB1 (MDR1) genetic variants are associated with methadone doses required for effective treatment of heroin dependence. *Hum Mol Genet.* 2008;17:2219–27.
67. Beyer A, Koch T, Schroder H, Schulz S, Holtt V. Effect of the A118G polymorphism on binding affinity, potency and agonist-mediated endocytosis, desensitization, and resensitization of the human mu-opioid receptor. *J Neurochem.* 2004;89:553–60.
68. Margas W, Zubkoff I, Schuler HG, Janicki PK, Ruiz-Velasco V. Modulation of Ca<sup>2+</sup> channels by heterologously expressed wild-type and mutant human micro-opioid receptors (hMORs) containing the A118G single-nucleotide polymorphism. *J Neurophysiol.* 2007;97:1058–67.
69. Krosiak T, Laforge KS, Gianotti RJ, Ho A, Nielsen DA, Kreek MJ. The single nucleotide polymorphism A118G alters functional properties of the human mu opioid receptor. *J Neurochem.* 2007;103:77–87.
70. Tan EC, Lim EC, Teo YY, Lim Y, Law HY, Sia AT. Ethnicity and OPRM1 variant independently predict pain perception and patient-controlled analgesia usage for post-operative pain. *Mol Pain.* 2009;5:32.
71. Walter C, Lotsch J. Meta-analysis of the relevance of the OPRM1 118A>G genetic variant for pain treatment. *Pain.* 2009;146:270–5.
72. Bruehl S, Chung OY, Donahue BS, Burns JW. Anger regulation style, postoperative pain, and relationship to the A118G mu opioid receptor gene polymorphism: a preliminary study. *J Behav Med.* 2006;29:161–9.
73. Chou WY, Wang CH, Liu PH, Liu CC, Tseng CC, Jawan B. Human opioid receptor A118G polymorphism affects intravenous patient-controlled analgesia morphine consumption after total abdominal hysterectomy. *Anesthesiology.* 2006;105:334–7.
74. Chou WY, Yang LC, Lu HF, et al. Association of mu-opioid receptor gene polymorphism (A118G) with variations in morphine consumption for analgesia after total knee arthroplasty. *Acta Anaesthesiol Scand.* 2006;50:787–92.
75. Fillingim RB, Kaplan L, Staud R, et al. The A118G single nucleotide polymorphism of the mu-opioid receptor gene (OPRM1) is associated with pressure pain sensitivity in humans. *J Pain.* 2005;6:159–67.
76. Klepstad P, Rakvag TT, Kaasa S, et al. The 118 A>G polymorphism in the human micro-opioid receptor gene may increase morphine requirements in patients with pain caused by malignant disease. *Acta Anaesthesiol Scand.* 2004;48:1232–9.
77. Oertel BG, Schmidt R, Schneider A, Geisslinger G, Lotsch J. The mu-opioid receptor gene polymorphism 118A>G depletes alfentanil-induced analgesia and protects against respiratory depression in homozygous carriers. *Pharmacogenet Genomics.* 2006;16:625–36.
78. Sia AT, Lim Y, Lim EC, et al. A118G single nucleotide polymorphism of human mu-opioid receptor gene influences pain perception and patient-controlled intravenous morphine consumption after

- intrathecal morphine for postcesarean analgesia. *Anesthesiology*. 2008;109:520–6.
79. Wand GS, McCaul M, Yang X, et al. The mu-opioid receptor gene polymorphism (A118G) alters HPA axis activation induced by opioid receptor blockade. *Neuropsychopharmacology*. 2002; 26:106–14.
80. Ginosar Y, Davidson EM, Meroz Y, Blotnick S, Shacham M, Caraco Y. Mu-opioid receptor (A118G) single-nucleotide polymorphism affects alfentanil requirements for extracorporeal shock wave lithotripsy: a pharmacokinetic-pharmacodynamic study. *Br J Anaesth*. 2009;103:420–7.
81. Wu WD, Wang Y, Fang YM, Zhou HY. Polymorphism of the micro-opioid receptor gene (OPRM1 118A>G) affects fentanyl-induced analgesia during anesthesia and recovery. *Mol Diagn Ther*. 2009;13:331–7.
82. Lotsch J, Stuck B, Hummel T. The human mu-opioid receptor gene polymorphism 118A>G decreases cortical activation in response to specific nociceptive stimulation. *Behav Neurosci*. 2006; 120:1218–24.
83. Lotsch J, Freynhagen R, Geisslinger G. Are polymorphisms in the mu-opioid receptor important for opioid therapy? *Schmerz*. 2005;19:378–82. 384–95.
84. Lotsch J, Geisslinger G. Relevance of frequent mu-opioid receptor polymorphisms for opioid activity in healthy volunteers. *Pharmacogenomics J*. 2006;6:200–10.
85. Lotsch J, Geisslinger G. Current evidence for a genetic modulation of the response to analgesics. *Pain*. 2006;121:1–5.
86. Lotsch J, Skarke C, Grosch S, Darimont J, Schmidt H, Geisslinger G. The polymorphism A118G of the human mu-opioid receptor gene decreases the pupil constrictory effect of morphine-6-glucuronide but not that of morphine. *Pharmacogenetics*. 2002;12:3–9.
87. Lotsch J, Zimmermann M, Darimont J, et al. Does the A118G polymorphism at the mu-opioid receptor gene protect against morphine-6-glucuronide toxicity? *Anesthesiology*. 2002;97:814–9.
88. Coulbault L, Beaussier M, Verstuft C, et al. Environmental and genetic factors associated with morphine response in the postoperative period. *Clin Pharmacol Ther*. 2006;79:316–24.
89. Fukuda K, Hayashida M, Ide S, et al. Association between OPRM1 gene polymorphisms and fentanyl sensitivity in patients undergoing painful cosmetic surgery. *Pain*. 2009;147:194–201.
90. Huehne K, Leis S, Muenster T, et al. High post surgical opioid requirements in Crohn's disease are not due to a general change in pain sensitivity. *Eur J Pain*. 2009;13:1036–42.
91. Janicki PK, Schuler G, Francis D, et al. A genetic association study of the functional A118G polymorphism of the human mu-opioid receptor gene in patients with acute and chronic pain. *Anesth Analg*. 2006;103:1011–7.
92. Hayashida M, Nagashima M, Satoh Y, et al. Analgesic requirements after major abdominal surgery are associated with OPRM1 gene polymorphism genotype and haplotype. *Pharmacogenomics*. 2008;9:1605–16.
93. Landau R, Kern C, Columb MO, Smiley RM, Blouin JL. Genetic variability of the mu-opioid receptor influences intrathecal fentanyl analgesia requirements in laboring women. *Pain*. 2008;139:5–14.
94. Reyes-Gibby CC, Shete S, Rakvag T, et al. Exploring joint effects of genes and the clinical efficacy of morphine for cancer pain: OPRM1 and COMT gene. *Pain*. 2007;130:25–30.
95. Ramchandani VA, Umhau J, Pavon FJ, et al. A genetic determinant of the striatal dopamine response to alcohol in men. *Mol Psychiatry*. 2011;16:809–17. Epub 2010 May 18.
96. Shabalina SA, Zaykin DV, Gris P, et al. Expansion of the human mu-opioid receptor gene architecture: novel functional variants. *Hum Mol Genet*. 2009;18:1037–51.
97. Berthele A, Platzer S, Jochim B, et al. COMT Val108/158Met genotype affects the mu-opioid receptor system in the human brain: evidence from ligand-binding, G-protein activation and preproenkephalin mRNA expression. *Neuroimage*. 2005;28:185–93.
98. Zubieta JK, Heitzeg MM, Smith YR, et al. COMT val158met genotype affects mu-opioid neurotransmitter responses to a pain stressor. *Science*. 2003;299:1240–3.
99. Rakvag TT, Klepstad P, Baar C, et al. The Val158Met polymorphism of the human catechol-O-methyltransferase (COMT) gene may influence morphine requirements in cancer pain patients. *Pain*. 2005;116:73–8.
100. Rakvag TT, Ross JR, Sato H, Skorpen F, Kaasa S, Klepstad P. Genetic variation in the catechol-O-methyltransferase (COMT) gene and morphine requirements in cancer patients with pain. *Mol Pain*. 2008;4:64.
101. Ross JR, Riley J, Taegetmeyer AB, et al. Genetic variation and response to morphine in cancer patients: catechol-O-methyltransferase and multidrug resistance-1 gene polymorphisms are associated with central side effects. *Cancer*. 2008;112:1390–403.
102. Nackley AG, Shabalina SA, Lambert JE, et al. Low enzymatic activity haplotypes of the human catechol-O-methyltransferase gene: enrichment for marker SNPs. *PLoS One*. 2009;4:e5237.
103. Nackley AG, Shabalina SA, Tchivileva IE, et al. Human catechol-O-methyltransferase haplotypes modulate protein expression by altering mRNA secondary structure. *Science*. 2006;314:1930–3.
104. Mogil JS, Ritchie J, Smith SB, et al. Melanocortin-1 receptor gene variants affect pain and mu-opioid analgesia in mice and humans. *J Med Genet*. 2005;42:583–7.
105. Mogil JS, Wilson SG, Chesler EJ, et al. The melanocortin-1 receptor gene mediates female-specific mechanisms of analgesia in mice and humans. *Proc Natl Acad Sci USA*. 2003;100:4867–72.
106. Esser R, Berry C, Du Z, et al. Preclinical pharmacology of lumiracoxib: a novel selective inhibitor of cyclooxygenase-2. *Br J Pharmacol*. 2005;144:538–50.
107. Cipollone F, Patrono C. Cyclooxygenase-2 polymorphism: putting a brake on the inflammatory response to vascular injury? *Arterioscler Thromb Vasc Biol*. 2002;22:1516–8.
108. Papafili A, Hill MR, Brull DJ, et al. Common promoter variant in cyclooxygenase-2 represses gene expression: evidence of role in acute-phase inflammatory response. *Arterioscler Thromb Vasc Biol*. 2002;22:1631–6.
109. Skarke C, Reus M, Schmidt R, et al. The cyclooxygenase 2 genetic variant -765 G>C does not modulate the effects of celecoxib on prostaglandin E2 production. *Clin Pharmacol Ther*. 2006;80:621–32.
110. Lee YS, Kim H, Wu TX, Wang XM, Dionne RA. Genetically mediated interindividual variation in analgesic responses to cyclooxygenase inhibitory drugs. *Clin Pharmacol Ther*. 2006;79:407–18.
111. Lotsch J, Fluhr K, Neddermayer T, Doehring A, Geisslinger G. The consequence of concomitantly present functional genetic variants for the identification of functional genotype-phenotype associations in pain. *Clin Pharmacol Ther*. 2009;85:25–30.
112. Kim H, Ramsay E, Lee H, Wahl S, Dionne RA. Genome-wide association study of acute post-surgical pain in humans. *Pharmacogenomics*. 2009;10:171–9.
113. Lotsch J, Geisslinger G. A critical appraisal of human genotyping for pain therapy. *Trends Pharmacol Sci*. 2010;31:312–7.



<http://www.springer.com/978-1-4614-1559-6>

Comprehensive Treatment of Chronic Pain by Medical,  
Interventional, and Integrative Approaches  
The AMERICAN ACADEMY OF PAIN MEDICINE Textbook on  
Patient Management

Deer, T.R.; Leong, M.S.; Buvanendran, A.; Gordin, V.;  
Kim, P.S.; Panchal, S.J.; Ray, A.L. (Eds.)

2013, XXVII, 1104 p. 412 illus., 233 illus. in color.,

Hardcover

ISBN: 978-1-4614-1559-6