
Methods of Cryopreservation in Fungi

2

Ladislav Homolka

Abstract

Traditional method of the routine subculturing by transfer of fungal cultures from staled to fresh media is not a very practical means of storing large numbers of fungal cultures. It is time-consuming, prone to contamination, and does not prevent genetic and physiological changes. At present, besides freeze-drying (lyophilization), cryopreservation seems to be the best preservation technique available for fungi.

Keywords

Cryopreservation • Fungi collections • Liquid nitrogen • Perlite
• Subculturing • Fungi storage

Introduction

Serious mycological (and generally biological) work requires a reliable source of cultures (i.e., well-defined and taxonomically determined starting material), which is ensured by its safe long-term storage. This implies the fundamental and growing importance of culture collections not only for preservation of the endangered genofond (and consequently the biodiversity), but also as a principal source of material for biotechnological processes, research, and teaching. The

first and most important problem to be solved is the long-term maintenance of this material.

Collections of fungi were originally kept by serial transfers from staled to fresh media. This routine subculturing is not a very practical method for storing large numbers of fungal cultures. It is time-consuming, prone to contamination, and does not prevent genetic and physiological changes (degeneration, aging) during long-term and frequent subculturing [1]. Over the years, various storage methods have been developed in order to eliminate these disadvantages. Their common feature is at least partial suppression of growth and metabolism of the cultures. Among them, keeping fungal cultures in sterile water [2–8] was surprisingly efficient (especially with lower fungi, but also with some basidiomycetes) and experiences its revival. In some fungi, preservation under a layer of mineral oil, in silica gel, soil, or sand [9–13] was successful. These

L. Homolka (✉)
Department of Ecology of Microorganisms,
Institute of Microbiology, Academy
of Sciences of the Czech Republic, Vídeňská 1083,
Prague 4 142 20, Czech Republic
e-mail: homolka@biomed.cas.cz

methods enabled a reduction in fungal growth and extended the time intervals between transfers to fresh media. Nevertheless, the method of serial subculturing is still used in collections with limited financial support or in the majority of other collections as a backup preservation method.

Searching for improved or new methods resulted through different intermediate steps in the introduction of lyophilization and cryopreservation of fungal cultures [14]. Application of the most extended method of culture preservation—freeze-drying (lyophilization), tested for sporulating fungi as early as 1945 [15]—is rather limited in the case of basidiomycetes and other fungi nonsporulating *in vitro* [16, 17]. Most attempts to revitalize dehydrated hyphae of fungi have failed, except for some successes [18–20]; nevertheless, the real absence of spores must be always carefully checked. Despite this, several attempts have recently been made using modified protocols [21–23] and the growing interest in this technique can be seen at present. An important role played in the whole process, besides the freeze-drying, is also the freezing rate and the lyoprotectant used [24, 25]. Freeze-drying of fungi has several important advantages over all other maintenance methods. Cultures can be stored easily in dense packing without any special requirements and need not be revived on agar slants prior to dispatch. The product is light, inactive, and dry, enabling easy distribution by mail.

The new or modified methods have been frequently used and evaluated [5, 6, 10, 26–31], but they are not generally applicable. Often a specific preservation protocol is necessary even for individual strains of the same species. At present, besides freeze-drying (lyophilization), cryopreservation seems to be the best preservation technique available for filamentous fungi [14, 26, 32]. A very comprehensive and detailed overview of the methods and results of the cryopreservation in microorganisms was published by Hubálek [33].

Cryogenic technique for long-term storage of large numbers of fungal species was introduced to ATCC in 1960 and the results have been very satisfactory [34, 35]. The technique was consecutively introduced to many other prominent

collections, e.g., CAB International Mycological Institute [1], etc. In certain collections—e.g., IFO (Institute for Fermentation Osaka)—nonsporulating cultures of basidiomycetes are stored by cryopreservation at -80°C in electric freezers [36].

Mycelium and/or spore suspensions with or without a cryoprotectant in sealed glass ampoules were originally used for cryopreservation of filamentous fungi. Later, glass ampoules were replaced with safer polypropylene cryovials and/or straws. Agar blocks immersed in an appropriate cryoprotectant were originally used as carriers of fungal mycelium for the cryopreservation process [37]. A useful straw technique with agar miniblocks for the preservation of fungi in liquid nitrogen was developed by Elliott in 1976 [38] and improved by Stalpers et al. [39]. Another technique using straws in cryotubes without a cryoprotectant solution was described by Hoffmann [40]. A modified Hoffmann's technique was compared with the original agar block one in our paper [29]. Commercial preservation systems with polystyrene beads as carriers were used for cryopreservation of conidia of entomopathogenic fungi [41] and of sporulating *Aspergillus fumigatus* cultures at -80°C [42]. Porous ceramic beads were employed for cryopreservation of several sporulating fungal cultures and for a *Saccharomyces cerevisiae* culture at -70°C by Palágyi et al. [43]. It is symptomatic that these techniques have not been used for nonsporulating filamentous fungi. In this context it should be mentioned that as early as 1978 Feltham et al. described a method of preservation of bacteria on glass beads at -76°C [44]. Some reports [45–47] indicate that cryopreservation at -80°C is suitable for many fungal cultures, including basidiomycetes. Nevertheless, Leeson et al. [48] state that to completely stabilize frozen cultures, the temperature must be sufficiently reduced to both minimize metabolism and prevent ice crystal formation, which can cause physical damage during storage. The temperature limit securing prevention of formation of such ice crystals is -139°C . This is why at present many culture collections start to keep their cultures at -150°C

in ultralow-temperature electric freezers, which are sometimes equipped with liquid nitrogen supply.

The cryopreservation process includes freezing and thawing and the protocol of these procedures plays an important role [49, 50]. In principle, there are two kinds of freezing protocols: a slow (controlled) one and a fast (uncontrolled) one, which both have been used for cryopreservation of fungi [51, 52]. Generally, too low freezing rates cause excessive dehydration and concentration of the solution leading to cell damage; on the contrary, too fast freezing leads to insufficient dehydration and formation of abundant ice crystals with lethal consequences. Nevertheless, different fungal cultures exhibit different sensitivities to freezing conditions and to the presence and concentration of cryoprotectants. A freezing rate of 1°C per minute is usually used for cryopreservation of fungi; in the author's experiments with sensitive mutant strains of *Agaricus bisporus*, the freezing rate 0.5°C per minute was successfully used.¹ Lately, cryomicroscopic methods have been used to study the process of freezing and thawing of fungal cultures [14, 24, 53]. Successful cryopreservation depends on the cryoprotectant used [54]. At present, dimethylsulfoxid and glycerol are the most widespread [55]. The method of cryogenic culture maintenance seems to be mostly successful also in nonsporulating cultures [14, 32, 56, 57]. An overview study on the influence of the cryopreservation process on survival of taxonomically very broad spectrum of fungi published by Smith [14] showed that there was no obvious link between taxonomic grouping and the response of the fungi to freezing and thawing. This was confirmed in our study [31]. Similar studies were carried out also in edible mushrooms *Lentinus edodes* [58] and the genus *Pleurotus* [59]. Davell and Flygh [60] showed that even an ectomycorrhizal fungus *Cantharellus cibarius* can be successfully cryopreserved when a sufficient number of cryoprotocols is tested. Cryopreservation of spores of

vesicular–arbuscular mycorrhizal fungi was described by Douds and Schenck [61].

Beyond survival, another principal requirement for the successful preservation of fungal strains is maintenance of their genetic and physiological features, such as growth, morphology, and metabolite production. In our experiments with some white-rot basidiomycetes, no negative effect of cryopreservation or the used cryoprotective on production of ligninolytic enzymes was found [62]. The complete revival of cryopreserved cultures (evaluated mostly by measuring the colony diameter) is generally still uncertain. The survival rate varies between 60 and 100% [21, 58, 59, 63, 64]. Only a few studies of the genetic stability of cryopreserved fungi have been performed. Singh et al. [47] confirmed the genetic stability of 11 cryogenically preserved edible mushroom strains by comparing random amplified polymorphic DNA (RAPD) and internal transcribed spacer (ITS) profiles. Using polymerase chain reaction (PCR) fingerprinting, Ryan et al. [65] checked the genetic stability of several isolates of *Fusarium oxysporum* and *Metarhizium anisopliae*. Other studies include confirmation of the genetic stability of *Uncinula necator* conidia after storage at –80°C [45] and investigation of the influence of mid-term cryopreservation at –80°C on 15 isolates of 10 basidiomycete species, for which the DNA fingerprint patterns were unchanged [66]. All of these reports were solely based on fingerprinting methods, which are not suitable for the detection of minute yet important changes in the genome, such as point or short indel mutations. Rather, sequencing approaches are required to successfully detect these mutations. This approach was used in our recent study [67]. Considering the above data, there is a continuous need for developing, improving, optimizing, and combining of preservation procedures, because the present methods are not applicable to all fungal cultures. Although many of these fungi can be grown in pure cultures on solid media, their growth is often attenuated and their morphology and other characteristics changed, which can result in their complete loss. The number of characteristics evaluating the success of preservation should be increased.

¹ Homolka, unpublished results.

As mentioned previously, cryopreservation, namely in liquid nitrogen, seems to be the most reliable, safe, and prospective method of a long-term maintenance of most fungal species, especially those not amenable to freeze-drying. It is probably the only storage technique that can ensure genomic and phenotypic stability. But not even the aforementioned cryopreservation method is applicable to preservation of all fungal cultures in the present form. According to the literature as well as the author's personal experience, especially the maintenance of basidiomycetes is challenging. Many of these fungi do not form asexual spores, their dominant life form, the vegetative mycelium, is sensitive to environmental conditions and therefore not amenable to freeze-drying.

To address these issues, a method of cryopreservation using perlite as a carrier for fungal mycelia was developed in the author's laboratory (perlite protocol or PP) [28] and then successfully verified for 442 basidiomycete strains [30]. Perlite is a unique aluminosilicate volcanic mineral that retains substantial amounts of water that can be released when needed—a feature that seems to have a dominant effect on cryopreservation success. The PP can be used for cryopreservation of taxonomically different groups of fungi, including yeasts [31], and works relatively well for fungi that cannot survive other routine preservation procedures. Expanded perlite was used as a solid support in solid-state fermentations [68]; otherwise it is used in many applications, particularly in the construction, horticulture, and other various industrial fields. It is recommended as an efficient purifying agent and as a carrier for pesticides, feed concentrates, herbicides, and other similar applications.

Perlite Protocol (PP)

The protocol is suitable for maintaining a broad spectrum of fungal cultures of different origin. It was verified in several culture collections (e.g., in Finland, the Netherlands, USA, Czech Republic, etc.) with great success and now it is routinely used there.

Materials

1. Distilled water.
2. Agar Difco.
3. Glycerol p.a. Sigma.
4. Isopropyl alcohol 100% Sigma (alternatively).
5. Agricultural-grade perlite (Agroperlite, GrowMarket s.r.o., Prague, Czech Republic, <http://www.growmarket.cz/produkt/agroperlite-81>)—particles 1–2 mm.
6. Dried wort extract Sladovit, Malthouse Bruntál, Bruntál, Czech Republic, diluted to a density of 4° Balling scale with distilled water (further wort in the text); or preferably MYA medium: Malt-extract Difco 25 g, Yeast-extract Difco 2 g, Glycerol p.a., Serva 50 g, distilled water ad 1,000 mL. pH adjusted to 6.5 with 1 M KOH solution.
7. Ethanol (70%).
8. Liquid nitrogen (LN).
9. Nunc CryoTube Vials 1.8 mL screw-capped (Nalgene/Nunc, Rochester, USA).
10. Cork borer.
11. Lancet and small spoon.
12. Rule for measuring of colony diameter.
13. Water bath.
14. Balling hydrometer (saccharometer) (alternatively).
15. Hot-air sterilizer.
16. Autoclave.
17. Refrigerator (about 4°C).
18. Deep-freezer (–80°C).
19. Microscope equipped with phase contrast.
20. Laminar flow box.
21. Thermostat (incubator) for 24°C.
22. Container for storing samples in liquid nitrogen (e.g., HARSCO TW-5K container, Harsco, Camp Hill, USA).
23. Programmable freezer for controlled freezing of cryovials with mycelia (e.g., IceCube 1800 freezer, SY-LAB Geraete GmbH, Neupurkersdorf, Austria); or alternatively Cryo 1°C Freezing Container “Mr. Frosty” (Nalgene Labware)².

² All chemicals and devices named can be replaced with other ones produced by other renowned companies.

Methods

The methods given below describe general procedures for cryopreservation of fungi using the perlite protocol (PP).

Strains (Cultures)

The starting cultures are kept on wort agar slants or dishes (wort 4° Balling, 1.5% agar Difco) at 4°C or other media suitable for the growth of the strains destined for the procedure (e.g., MYA medium with 1.5% agar, etc.) and transferred to the fresh medium every 6 months.

1. Prepare an agar medium, sterilize it in an autoclave (121°C, 20 min.), pour it into sterile plastic Petri dishes (diameter 100 mm, 30 mL per dish), and let it cool down in a laminar flow box.
2. In a laminar flow box cut out an agar plug (6 mm diameter) from the actively growing part of a colony on a Petri dish with a cork borer, place it on a Petri dish with fresh medium using a sterile lancet and then let the dish incubate for 14 days at 24°C. Then put the dish(es) into a refrigerator and keep it at about 4°C.

Culture Preparation and Freezing-Thawing Protocols

Fungal cultures are grown directly in firmly closed sterile plastic cryovials (1.8 mL) with 200 mg of perlite (Agroperlite, agricultural grade) moistened with 1 mL of wort (4° Balling) or other medium (e.g., MYA) enriched with 5% glycerol as a cryoprotectant. For sterilization of the cork borer, lancet, and small spoon use a hot-air sterilizer (150°C, 30 min.).

1. Distribute perlite into cryovials (200 mg per vial), flood it with 1.8 mL of the medium enriched with 5% of glycerol, and sterilize vials in autoclave (121°C, 20 min.). In a laminar flow box cut out an agar plug (6 mm diameter) from the actively growing part of a colony

on a Petri dish using a cork borer, place it using a sterile lancet on the surface of perlite in the cryovial, close the vial firmly and let it incubate for 14 days at 24°C.

2. Freeze the cryovials with perlite overgrown by mycelium in a programmable freezer (or alternatively in a “Mr. Frosty” container in a deep-freezer) to –70°C at a freezing rate of 1°C per minute. Then place them in LN in a container.
3. Take the stored frozen cultures in cryovials out of the LN container, transfer them to a water bath (37°C), and leave them there until the ice is completely thawed (thawing—reactivation of cultures). Prior to opening, disinfect the surface of cryovials with 70% ethanol.

Viability Test

1. After thawing, separate at least partially the perlite particles overgrown with mycelium by shaking, the content of the cryovials (two parallels of each strain) divide into three approximately equal aliquots each and these plate onto wort (or an other) agar medium in Petri dishes (diameter 100 mm) using a small sterile spoon.
2. Incubate the cultures in Petri dishes at 24°C for 14 days. Strains exhibiting survival of at least four out of six separate aliquots are considered viable.

Growth Estimation and Morphological Analysis

1. Growth of cultures measure as a mean diameter increase of a growth-covered zone (in mm) during a 14-day incubation at 24°C on the respective agar medium in Petri dishes (diameter 100 mm) inoculated with perlite aliquots from cryovials before freezing and after reactivation. Measure six zones (three aliquots from two cryovials) for each strain. The first occurrence of growth varies between frozen cultures, with some strains showing signs of re-growth within 2 days but most strains reactivating within 7 days after plating.

2. Use the same procedure except for freezing and thawing for growth measurement of the control.
3. Carry out the morphological analysis on control cultures and on those arising from the viability tests. Check the selected macroscopic features (colony color, reverse color, texture of the mycelium) and microscopic features (hyphal branching, presence/absence of clamp connections, presence/absence of hyphal vacuolization, etc.) using a microscope.
4. If possible, estimate also other characteristics of the resulting cultures (e.g., enzyme or metabolite production, etc.) according to your consideration.

References

1. Onions AHS (1971) Preservation of fungi. In: Booth C (ed) *Methods in microbiology*, vol 4. Academic, New York, London, pp 113–115
2. Marx DH, Daniel WJ (1976) Maintaining cultures of ectomycorrhizal and plant pathogenic fungi in sterile water cold storage. *Can J Microbiol* 22:338–341
3. Ellis JJ (1979) Preserving fungus strains in sterile water. *Mycologia* 71:1072–1075
4. Richter DL, Bruhn JN (1989) Revival of saprotrophic and mycorrhizal basidiomycete cultures from cold storage in sterile water. *Can J Microbiol* 35: 1055–1060
5. Smith JE, McKay D, Molina R (1994) Survival of mycorrhizal fungal isolates stored in sterile water at two temperatures and retrieved on solid and liquid nutrient media. *Can J Microbiol* 40:736–742
6. Burdsall HH, Dorworth EB (1994) Preserving cultures of wood-decaying Basidiomycotina using sterile distilled water in cryovials. *Mycologia* 86:275–280
7. Borman AM, Szekely A, Campbell CK, Johnson EM (2006) Evaluation of the viability of pathogenic filamentous fungi after prolonged storage in sterile water and review of recent published studies on storage methods. *Mycopathologia* 161:361–368
8. Richter DL (2008) Revival of saprotrophic and mycorrhizal basidiomycete cultures after 20 years in cold storage in sterile water. *Can J Microbiol* 54:595–599
9. Perrin PW (1979) Long-term storage of cultures of wood-inhabiting fungi under mineral oil. *Mycologia* 71:867–869
10. Johnson GC, Martin AK (1992) Survival of wood-inhabiting fungi stored for 10 years in water and under oil. *Can J Microbiol* 38:861–864
11. Sharma B, Smith D (1999) Recovery of fungi after storage for over a quarter of a century. *World J Microbiol Biotechnol* 15:517–519
12. Delcán J, Moyano C, Raposo R, Melgarejo P (2002) Storage of *Botrytis cinerea* using different methods. *J Plant Pathol* 84:3–9
13. Baskarathevan J, Jaspers MV, Jones EE, Ridgway HJ (2009) Evaluation of different storage methods for rapid and cost-effective preservation *Botryosphaera* species. *N Z Plant Protect* 62:234–237
14. Smith D (1998) The use of cryopreservation in the ex-situ conservation of fungi. *CryoLetters* 19:79–90
15. Raper KB, Alexander DF (1945) Preservation of molds by lyophil process. *Mycologia* 37:499–525
16. Antheunisse J (1973) Viability of lyophilized microorganisms after storage. *Antonie Leeuwenhoek* 39: 243–248
17. Hwang S-W, Kwolek WF, Haynes WC (1976) Investigation of ultra-low temperature for fungal cultures. III. Viability and growth rate of mycelial cultures following cryogenic storage. *Mycologia* 68:377–387
18. Bazzigher G (1962) Ein vereinfachtes Gefriertrocknungsverfahren zur Konservierung von Pilzkulturen. *Phytopathol Z* 45:53–56
19. Pertot E, Puc A, Kremser M (1977) Lyophilization of nonsporulating strains of the fungus *Claviceps*. *Eur J Appl Microbiol* 4:289–294
20. Tommerup IC (1988) Long-term preservation by L-drying and storage of vesicular arbuscular mycorrhizal fungi. *Trans Brit Mycol Soc* 90:585–591
21. Tan CS, Stalpers JA, van Ingen CW (1991) Freeze-drying of fungal hyphae. *Mycologia* 83:654–657
22. Croan SC, Burdsall HH Jr, Rentmeester RM (1999) Preservation of tropical wood-inhabiting basidiomycetes. *Mycologia* 91:908–916
23. Sundari SK, Adholeya A (1999) Freeze-drying vegetative mycelium of *Laccaria fraterna* and its subsequent regeneration. *Biotechnol Tech* 13:491–495
24. Tan CS, Vlug IJA, Stalpers JA, van Ingen CW (1994) Microscopical observations on the influence of the cooling rate during freeze-drying of conidia. *Mycologia* 86:281–289
25. Tan CS, van Ingen CW, Talsma H, van Miltenburg JC, Steffensen CL, Vlug IJA, Stalpers JA (1995) Freeze-drying of fungi, influence of composition and glass transition temperature of the protectant. *Cryobiology* 32:60–67
26. Ryan MJ, Smith D (2004) Fungal genetic resource centres and the genomic challenge. *Mycol Res* 108:1351–1362
27. Ryan MJ, Smith D (2007) Cryopreservation and freeze-drying of fungi employing centrifugal and shelf freeze-drying. *Meth Mol Biol* 368:127–140
28. Homolka L, Lisá L, Eichlerová I, Nerud F (2001) Cryopreservation of basidiomycete strains using perlite. *J Microbiol Meth* 47:307–313
29. Homolka L, Lisá L, Nerud F (2003) Viability of basidiomycete strains after cryopreservation, comparison of two different freezing protocols. *Folia Microbiol* 48:219–226
30. Homolka L, Lisá L, Nerud F (2006) Basidiomycete cryopreservation on perlite, evaluation of a new method. *Cryobiology* 52:446–453

31. Homolka L, Lisá L, Kubátová A, Váňová M, Janderová B, Nerud F (2007) Cryopreservation of filamentous micromycetes and yeasts using perlite. *Folia Microbiol* 52:153–157
32. Challen MP, Elliott T (1986) Polypropylene straw ampoules for the storage of microorganisms in liquid nitrogen. *J Microbiol Meth* 5:11–23
33. Hubálek Z (1996) Cryopreservation of microorganisms. Academia Publishing House, Prague
34. Hwang S-W (1960) Effects of ultra-low temperatures on the viability of selected fungus strains. *Mycologia* 52:527–529
35. Smith D (1983) Cryoprotectants and the cryopreservation of fungi. *Trans Brit Mycol Soc* 80:360–363
36. Ito T (1996) Preservation of fungal cultures at the Institute of Fermentation, Osaka (IFO). In: Samson RA, Stalpers JA, van der Mei D, Stouthamer AH (eds) Culture collections to improve the quality of life. The Netherlands and the World Federation for Culture Collections, Centraalbureau voor Schimmelcultures, Baarn, pp 210–211
37. Hwang S-W (1968) Investigation of ultra-low temperature for fungal cultures. I. An evaluation of liquid nitrogen storage for preservation of selected fungal cultures. *Mycologia* 60:613–621
38. Elliott TJ (1976) Alternative ampoule for storing fungal cultures in liquid nitrogen. *Trans Brit Mycol Soc* 67:545–546
39. Stalpers JA, de Hoog A, Vlug IJ (1987) Improvement of the straw technique for the preservation of fungi in liquid nitrogen. *Mycologia* 79:82–89
40. Hoffmann P (1991) Cryopreservation of fungi. *World J Microbiol Biotechnol* 7:92–94
41. Chandler D (1994) Cryopreservation of fungal spores using porous beads. *Mycol Res* 98:525–526
42. Belkacemi L, Barton RC, Evans EGV (1997) Cryopreservation of *Aspergillus fumigatus* stock cultures with a commercial bead system. *Mycoses* 40:103–104
43. Palágyi Z, Nagy Á, Vastag M, Ferency L, Vágvölgyi C (1997) Maintenance of fungal strains on cryopreservative-immersed porous ceramic beads. *Biotechnol Tech* 11:249–250
44. Feltham RKA, Power AK, Pell PA, Sneath PHA (1978) A simple method for storage of bacteria at -76° C. *J Appl Bacteriol* 44:313–316
45. Stummer BE, Zanker T, Scott ES (1999) Cryopreservation of airdried conidia of *Uncinula necator*. *Austr Plant Pathol* 28:82–84
46. Kitamoto Y, Suzuki A, Shimada S, Yamanaka K (2002) A new method for the preservation of fungus stock cultures by deepfreezing. *Mycoscience* 43:143–149
47. Singh SK, Upadhyay RC, Kamal S, Tiwari M (2004) Mushroom cryopreservation and its effect on survival, yield and genetic stability. *CryoLetters* 25:23–32
48. Leeson EA, Cann JP, Morris GJ (1984) Maintenance of algae and protozoa. In: Kirsop BE, Snell JJS (eds) Maintenance of microorganisms. Academic, London, pp 131–160
49. Leef J, Mazur P (1978) Physiological response of *Neurospora* conidia to freezing in the dehydrated, hydrated or germinated state. *Appl Environ Microbiol* 35:72–83
50. Morris GJ, Smith D, Coulson GE (1988) A comparative study of the changes in the morphology of hyphae during freezing and viability upon thawing for twenty species of fungi. *J Gen Microbiol* 134:2897–2906
51. Goos RD, Davis EE, Butterfield W (1967) Effect of warming rates on the viability of frozen fungus spores. *Mycologia* 59:58–66
52. Dahmen H, Staub T, Schwinn FT (1983) Technique for long-term preservation of phytopathogenic fungi in liquid nitrogen. *Phytopathology* 73:241–246
53. Smith D, Coulson GE, Morris GJ (1986) A comparative study of the morphology and viability of hyphae of *Penicillium expansum* and *Phytophthora nicotianae* during freezing and thawing. *J Gen Microbiol* 132:2013–2021
54. Jong SC, Davis EE (1978) Conservation of reference strains of *Fusarium* in pure culture. *Mycopathologia* 66:153–159
55. Hubálek Z (2003) Protectants used in the cryopreservation of microorganisms. *Cryobiology* 46:205–229
56. Homolka L (1976) On the problem of maintenance and cultivation of higher fungi. *Folia Microbiol* 21:189–190
57. Chvostová V, Nerud F, Homolka L (1995) Viability of wood-inhabiting basidiomycetes following cryogenic preservation. *Folia Microbiol* 40:193–197
58. Roquebert MF, Bury E (1993) Effect of freezing and thawing on cell membranes of *Lentinus edodes*, the Shiitake mushroom. *World J Microbiol Biotechnol* 9:641–647
59. Mata G, Salmones D, Pérez R, Guzmán G (1994) Behavior of some strains of the genus *Pleurotus* after different procedures for freezing in liquid nitrogen. *Rev Microbiol Sao Paulo* 25:197–200
60. Danell E, Flygh G (2002) Cryopreservation of the ectomycorrhizal mushroom *Cantharellus cibarius*. *Mycol Res* 106:1340–1342
61. Douds DD Jr, Schenck NC (1990) Cryopreservation of spores of vesicular-arbuscular mycorrhizal fungi. *New Phytol* 115:667–674
62. Stoychev I, Homolka L, Nerud F, Lisá L (1998) Activities of ligninolytic enzymes in some white-rot basidiomycete strains after recovering from cryopreservation in liquid nitrogen. *Antonie Leeuwenhoek* 73:211–214
63. Schipper MAA, Bekker-Holtman J (1976) Viability of lyophilized fungal cultures. *Antonie Leeuwenhoek* 42:325–328
64. Smith D, Ward SM (1987) Notes on the preservation of fungi. CAB International Mycological Institute, Slough, UK

-
65. Ryan MJ, Jeffries P, Bridge PD, Smith D (2001) Developing cryopreservation protocols to secure fungal gene function. *CryoLetters* 22:115–124
66. Voyron S, Roussel S, Munaut F, Varese GC, Ginepro M, Declerck S, Marchisio VF (2009) Vitality and genetic fidelity of white-rot fungi mycelia following different methods of preservation. *Mycol Res* 113:1027–1038
67. Homolka L, Lisá L, Eichlerová I, Valášková V, Baldrian P (2010) Effect of long-term preservation of different basidiomycetes on perlite in liquid nitrogen on their growth, morphological, enzymatic and genetic characteristics. *Fungal Biol* 114:929–935
68. Kerem Z, Hadar Y (1993) Effect of manganese on lignin degradation by *Pleurotus ostreatus* during solid-state fermentation. *Appl Environ Microbiol* 59:4115–4120



<http://www.springer.com/978-1-4614-2355-3>

Laboratory Protocols in Fungal Biology

Current Methods in Fungal Biology

Gupta, V.K.; Tuohy, M.G.; Ayyachamy, M.; Turner, K.M.;

O'Donovan, A. (Eds.)

2013, XXV, 604 p., Hardcover

ISBN: 978-1-4614-2355-3