

Bivariate Right Fractional Pseudo-Polynomial Monotone Approximation

George A. Anastassiou

Abstract In this article we deal with the following general two-dimensional problem: Let f be a two variable continuously differentiable real valued function of a given order, let \bar{L} be a linear right fractional mixed partial differential operator and suppose that $\bar{L}(f) \geq 0$ on a critical region. Then for sufficiently large $n, m \in \mathbb{N}$, we can find a sequence of pseudo-polynomials $Q_{n,m}^*$ in two variables with the property $\bar{L}(Q_{n,m}^*) \geq 0$ on this critical region such that f is approximated with rates right fractionally and simultaneously by $Q_{n,m}^*$ in the uniform norm on the whole domain of f . This restricted approximation is given via inequalities involving the mixed modulus of smoothness $\omega_{s,q}$, $s, q \in \mathbb{N}$, of highest order integer partial derivative of f .

1 Introduction

The topic of monotone approximation started in [10] has become a major trend in approximation theory. A typical problem in this subject is: given a positive integer k , approximate a given function whose k th derivative is ≥ 0 by polynomials having this property.

In [3] the authors replaced the k th derivative with a linear differential operator of order k . We mention this motivating result.

Theorem 1 *Let h, k, p be integers, $0 \leq h \leq k \leq p$ and let f be a real function, $f^{(p)}$ continuous in $[-1, 1]$ with modulus of continuity $\omega_1(f^{(p)}, x)$ there. Let $a_j(x)$, $j = h, h + 1, \dots, k$ be real functions, defined and bounded on $[-1, 1]$ and assume $a_h(x)$ is either \geq some number $\alpha > 0$ or \leq some number $\beta < 0$ throughout $[-1, 1]$. Consider the operator*

$$L = \sum_{j=h}^k a_j(x) \left[\frac{d^j}{dx^j} \right] \quad (1)$$

G.A. Anastassiou (✉)
Department of Mathematical Sciences, University of Memphis,
Memphis, TN 38152, USA
e-mail: ganastss@memphis.edu

and suppose, throughout $[-1, 1]$,

$$L(f) \geq 0. \quad (2)$$

Then, for every integer $n \geq 1$, there is a real polynomial $Q_n(x)$ of degree $\leq n$ such that

$$L(Q_n) \geq 0 \text{ throughout } [-1, 1] \quad (3)$$

and

$$\max_{-1 \leq x \leq 1} |f(x) - Q_n(x)| \leq Cn^{k-p} \omega_1 \left(f^{(p)}, \frac{1}{n} \right), \quad (4)$$

where C is independent of n or f .

Next let $n, m \in \mathbb{Z}_+$, P_θ denote the space of algebraic polynomials of degree $\leq \theta$. Consider the tensor product spaces $P_n \otimes C([-1, 1])$, $C([-1, 1]) \otimes P_m$ and their sum $P_n \otimes C([-1, 1]) + C([-1, 1]) \otimes P_m$, that is

$$\begin{aligned} & P_n \otimes C([-1, 1]) + C([-1, 1]) \otimes P_m \quad (5) \\ &= \left\{ \sum_{i=0}^n x^i A_i(y) + \sum_{j=0}^m B_j(x) y^j; A_i, B_j \in C([-1, 1]), x, y \in [-1, 1] \right\}. \end{aligned}$$

This is the space of pseudo-polynomials of degree $\leq (n, m)$, first introduced by A. Marchaud in 1924–1927 (see [7, 8]). Here $f^{(k,l)}$ denotes $\frac{\partial^{k+l} f}{\partial x^k \partial y^l}$, the (k, l) -partial derivative of f .

In this section we consider the space $C^{r,p}([-1, 1]^2) = \{f : [-1, 1]^2 \rightarrow \mathbb{R}; f^{(k,l)}$ is continuous for $0 \leq k \leq r, 0 \leq l \leq p\}$. Let $f \in C([-1, 1]^2)$; for $\delta_1, \delta_2 \geq 0$, define the mixed modulus of smoothness of order (s, q) , $s, q \in \mathbb{N}$ (see [9], pp. 516–517) by

$$\begin{aligned} \omega_{s,q}(f; \delta_1, \delta_2) &\equiv \sup \left\{ \left| {}_x \Delta_{h_1}^s \circ_y \Delta_{h_2}^q f(x, y) \right| : (x, y), \right. \\ &\quad \left. (x + sh_1, y + qh_2) \in [-1, 1]^2, |h_i| \leq \delta_i, i = 1, 2 \right\}. \quad (6) \end{aligned}$$

Here

$$\begin{aligned} & {}_x \Delta_{h_1}^s \circ_y \Delta_{h_2}^q f(x, y) \\ &\equiv \sum_{\sigma=0}^s \sum_{\mu=0}^q (-1)^{s+q-\sigma-\mu} \binom{s}{\sigma} \binom{q}{\mu} f(x + \sigma h_1, y + \mu h_2) \quad (7) \end{aligned}$$

is a mixed difference of order (s, q) .

We mention

Theorem 2 (see Gonska [4]) *Let $r, p \in \mathbb{Z}_+$, $s, q \in \mathbb{N}$, and $f \in C^{r,p}([-1, 1]^2)$. Let $n, m \in \mathbb{N}$ with $n \geq \max\{4(r+1), r+s\}$ and $m \geq \max\{4(p+1), p+q\}$. Then there exists a linear operator $Q_{n,m}$ from $C^{r,p}([-1, 1]^2)$ into the space of pseudopolynomials $(P_n \otimes C([-1, 1]) + C([-1, 1]) \otimes P_m)$ such that*

$$\begin{aligned} & \left| (f - Q_{n,m}(f))^{(k,l)}(x, y) \right| \\ & \leq M_{r,s} \cdot M_{p,q} (\Delta_n(x))^{r-k} \cdot (\Delta_m(y))^{p-l} \cdot \omega_{s,q}(f^{(r,p)}; \Delta_n(x), \Delta_m(y)), \end{aligned} \quad (8)$$

for all $(0, 0) \leq (k, l) \leq (r, p)$, $x, y \in [-1, 1]$, where

$$\Delta_\theta(z) = \frac{\sqrt{1-z^2}}{\theta} + \frac{1}{\theta^2}, \quad \theta = n, m; \quad z = x, y \in [-1, 1].$$

The constants $M_{r,s}, M_{p,q}$, are independent of $f, (x, y)$ and (n, m) ; they depend only on $(r, s), (p, q)$, respectively.

See also [5], saying that $Q_{n,m}^{(r,p)}(f)$ is continuous on $[-1, 1]^2$.

The need following result which is an easy consequence of the last theorem (see [9, p. 517]).

Corollary 3 *Let $r, p \in \mathbb{Z}_+$, $s, q \in \mathbb{N}$, and $f \in C^{r,p}([-1, 1]^2)$. Let $n, m \in \mathbb{N}$ with $n \geq \max\{4(r+1), r+s\}$ and $m \geq \max\{4(p+1), p+q\}$. Then there exists a pseudopolynomial*

$$Q_{n,m} \equiv Q_{n,m}(f) \in (P_n \otimes C([-1, 1]) + C([-1, 1]) \otimes P_m)$$

such that

$$\|f^{(k,l)} - Q_{n,m}^{(k,l)}\|_\infty \leq \frac{\dot{C}}{n^{r-k}m^{p-l}} \cdot \omega_{s,q}\left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m}\right), \quad (9)$$

for all $(0, 0) \leq (k, l) \leq (r, p)$. Here the constant \dot{C} depends only on r, p, s, q .

Corollary 3 was used in the proof of the main motivational result that follows.

Theorem 4 (see [1]) *Let h_1, h_2, v_1, v_2, r, p be integers, $0 \leq h_1 \leq v_1 \leq r$, $0 \leq h_2 \leq v_2 \leq p$ and let $f \in C^{r,p}([-1, 1]^2)$, with $f^{(r,p)}$ having a mixed modulus of smoothness $\omega_{s,q}(f^{(r,p)}; x, y)$ there, $s, q \in \mathbb{N}$. Let $\alpha_{i,j}(x, y)$, $i = h_1, h_1 + 1, \dots, v_1$; $j = h_2, h_2 + 1, \dots, v_2$ be real-valued functions, defined and bounded in $[-1, 1]^2$ and suppose $\alpha_{h_1 h_2}$ is either $\geq \alpha > 0$ or $\leq \beta < 0$ throughout $[-1, 1]^2$. Take the operator*

$$L = \sum_{i=h_1}^{v_1} \sum_{j=h_2}^{v_2} \alpha_{ij}(x, y) \frac{\partial^{i+j}}{\partial x^i \partial y^j} \quad (10)$$

and assume, throughout $[-1, 1]^2$ that

$$L(f) \geq 0. \quad (11)$$

Then for any integers n, m with $n \geq \max\{4(r+1), r+s\}$, $m \geq \max\{4(p+1), p+q\}$, there exists a pseudopolynomial

$$\mathcal{Q}_{n,m} \in (P_n \otimes C([-1, 1]) + C([-1, 1]) \otimes P_m)$$

such that $L(\mathcal{Q}_{m,n}) \geq 0$ throughout $[-1, 1]^2$ and

$$\|f^{(k,l)} - \mathcal{Q}_{n,m}^{(k,l)}\|_\infty \leq \frac{C}{n^{r-v_1} m^{p-v_2}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \quad (12)$$

for all $(0, 0) \leq (k, l) \leq (h_1, h_2)$. Moreover we get

$$\|f^{(k,l)} - \mathcal{Q}_{n,m}^{(k,l)}\|_\infty \leq \frac{C}{n^{r-k} m^{p-l}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \quad (13)$$

for all $(h_1 + 1, h_2 + 1) \leq (k, l) \leq (r, p)$. Also (13) is valid whenever $0 \leq k \leq h_1$, $h_2 + 1 \leq l \leq p$ or $h_1 + 1 \leq k \leq r$, $0 \leq l \leq h_2$. Here C is a constant independent of f and n, m . It depends only on r, p, s, q, L .

We are also motivated by [2].

We need

Definition 5 (see [6]) Let $[-1, 1]^2$; $\alpha_1, \alpha_2 > 0$; $\alpha = (\alpha_1, \alpha_2)$, $f \in C([-1, 1]^2)$, $x = (x_1, x_2)$, $t = (t_1, t_2) \in [-1, 1]^2$. We define the right mixed Riemann-Liouville fractional two dimensional integral of order α

$$\begin{aligned} & (I_{1-}^\alpha f)(x) \quad (14) \\ & := \frac{1}{\Gamma(\alpha_1) \Gamma(\alpha_2)} \int_{x_1}^1 \int_{x_2}^1 (t_1 - x_1)^{\alpha_1-1} (t_2 - x_2)^{\alpha_2-1} f(t_1, t_2) dt_1 dt_2, \end{aligned}$$

with $x_1, x_2 < 1$. Notice here that $I_{1-}^\alpha(|f|) < \infty$.

Definition 6 Let $\alpha_1, \alpha_2 > 0$ with $[\alpha_1] = m_1$, $[\alpha_2] = m_2$, ($[\cdot]$ ceiling of the number). Let here $f \in C^{m_1, m_2}([-1, 1]^2)$. We consider the right Caputo type fractional partial derivative:

$$\begin{aligned} & D_{1-}^{(\alpha_1, \alpha_2)} f(x) \\ & := \frac{(-1)^{m_1+m_2}}{\Gamma(m_1 - \alpha_1) \Gamma(m_2 - \alpha_2)} \quad (15) \end{aligned}$$

$$\cdot \int_{x_1}^1 \int_{x_2}^1 (t_1 - x_1)^{m_1 - \alpha_1 - 1} (t_2 - x_2)^{m_2 - \alpha_2 - 1} \frac{\partial^{m_1 + m_2} f(t_1, t_2)}{\partial t_1^{m_1} \partial t_2^{m_2}} dt_1 dt_2,$$

$\forall x = (x_1, x_2) \in [-1, 1]^2$, where Γ is the gamma function

$$\Gamma(v) = \int_0^\infty e^{-t} t^{v-1} dt, \quad v > 0. \tag{16}$$

We set

$$D_{1-}^{(0,0)} f(x) := f(x), \quad \forall x \in [-1, 1]^2; \tag{17}$$

$$D_{1-}^{(m_1, m_2)} f(x) := (-1)^{m_1 + m_2} \frac{\partial^{m_1 + m_2} f(x)}{\partial x_1^{m_1} \partial x_2^{m_2}}, \quad \forall x \in [-1, 1]^2. \tag{18}$$

Definition 7 We also set

$$D_{1-}^{(0, \alpha_2)} f(x) := \frac{(-1)^{m_2}}{\Gamma(m_2 - \alpha_2)} \int_{x_2}^1 (t_2 - x_2)^{m_2 - \alpha_2 - 1} \frac{\partial^{m_2} f(x_1, t_2)}{\partial t_2^{m_2}} dt_2, \tag{19}$$

$$D_{1-}^{(\alpha_1, 0)} f(x) := \frac{(-1)^{m_1}}{\Gamma(m_1 - \alpha_1)} \int_{x_1}^1 (t_1 - x_1)^{m_1 - \alpha_1 - 1} \frac{\partial^{m_1} f(t_1, x_2)}{\partial t_1^{m_1}} dt_1, \tag{20}$$

and

$$D_{1-}^{(m_1, \alpha_2)} f(x) := \frac{(-1)^{m_2}}{\Gamma(m_2 - \alpha_2)} \int_{x_2}^1 (t_2 - x_2)^{m_2 - \alpha_2 - 1} \frac{\partial^{m_1 + m_2} f(x_1, t_2)}{\partial x_1^{m_1} \partial t_2^{m_2}} dt_2, \tag{21}$$

$$D_{1-}^{(\alpha_1, m_2)} f(x) := \frac{(-1)^{m_1}}{\Gamma(m_1 - \alpha_1)} \int_{x_1}^1 (t_1 - x_1)^{m_1 - \alpha_1 - 1} \frac{\partial^{m_1 + m_2} f(t_1, x_2)}{\partial t_1^{m_1} \partial x_2^{m_2}} dt_1. \tag{22}$$

In this article we extend Theorem 4 to the fractional level. Indeed here L is replaced by \bar{L} , a linear right Caputo fractional mixed partial differential operator. Now the monotonicity property holds true only on the critical square of $[-1, 0]^2$. Simultaneously fractional convergence remains true on all of $[-1, 1]^2$.

2 Main Result

We present

Theorem 8 *Let h_1, h_2, v_1, v_2, r, p be integers, $0 \leq h_1 \leq v_1 \leq r, 0 \leq h_2 \leq v_2 \leq p$ and let $f \in C^{r,p}([-1, 1]^2)$, with $f^{(r,p)}$ having a mixed modulus of smoothness $\omega_{s,q}(f^{(r,p)}; x, y)$ there, $s, q \in \mathbb{N}$. Let $\alpha_{ij}(x, y), i = h_1, h_1 + 1, \dots, v_1; j = h_2, h_2 + 1, \dots, v_2$ be real valued functions, defined and bounded in $[-1, 1]^2$ and suppose $\alpha_{h_1 h_2}$ is either $\geq \alpha > 0$ or $\leq \beta < 0$ throughout $[-1, 0]^2$. Assume that $h_1 + h_2 = 2\gamma, \gamma \in \mathbb{Z}_+$. Here $n, m \in \mathbb{N} : n \geq \max\{4(r+1), r+s\}, m \geq \max\{4(p+1), p+q\}$. Set*

$$l_{ij} := \sup_{(x,y) \in [-1,1]^2} |\alpha_{h_1 h_2}^{-1}(x, y) \alpha_{ij}(x, y)| < \infty, \quad (23)$$

for all $h_1 \leq i \leq v_1, h_2 \leq j \leq v_2$. Let $\alpha_{1i}, \alpha_{2j} > 0, \alpha_{1i}, \alpha_{2j} \notin \mathbb{N}$, with $[\alpha_{1i}] = i, [\alpha_{2j}] = j, i = 0, 1, \dots, r; j = 0, 1, \dots, p, ([\cdot]$ ceiling of the number), $\alpha_{10} = 0, \alpha_{20} = 0$.

Consider the right fractional bivariate differential operator

$$\bar{L} := \sum_{i=h_1}^{v_1} \sum_{j=h_2}^{v_2} \alpha_{ij}(x, y) D_{1-}^{(\alpha_{1i}, \alpha_{2j})}. \quad (24)$$

Assume $\bar{L}f(x, y) \geq 0$, on $[-1, 0]^2$. Then there exists

$$Q_{n,m}^* \equiv Q_{n,m}^*(f) \in (P_n \otimes C([-1, 1]) + C([-1, 1]) \otimes P_m)$$

such that $\bar{L}Q_{n,m}^*(x, y) \geq 0$, on $[-1, 0]^2$. Furthermore it holds:

1.

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})}(f) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^* \right\|_{\infty, [-1, 1]^2} \\ & \leq \frac{C 2^{(i+j) - (\alpha_{1i} + \alpha_{2j})}}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1) n^{r-i} m^{p-j}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \end{aligned} \quad (25)$$

where C is a constant that depends only on $r, p, s, q; (h_1 + 1, h_2 + 1) \leq (i, j) \leq (r, p)$, or $0 \leq i \leq h_1, h_2 + 1 \leq j \leq p$, or $h_1 + 1 \leq i \leq r, 0 \leq j \leq h_2$,

2.

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})}(f) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^* \right\|_{\infty, [-1, 1]^2} \\ & \leq \frac{C_{ij}}{n^{r-v_1} m^{p-v_2}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \end{aligned} \quad (26)$$

for $(1, 1) \leq (i, j) \leq (h_1, h_2)$, where $c_{ij} = \dot{C} A_{ij}$, with

$$\begin{aligned}
 & A_{ij} \\
 & := \left\{ \left[\sum_{\tau=h_1}^{v_1} \sum_{\mu=h_2}^{v_2} \frac{l_{\tau\mu} 2^{(\tau+\mu)-(\alpha_{1\tau}+\alpha_{2\mu})}}{\Gamma(\tau - a_{1\tau} + 1) \Gamma(\mu - \alpha_{2\mu} + 1)} \right] \right. \\
 & \cdot \left(\sum_{k=0}^{h_1-i} \frac{2^{h_1-\alpha_{1i}-k}}{k! \Gamma(h_1 - \alpha_{1i} - k + 1)} \right) \\
 & \cdot \left. \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2 - \alpha_{2j} - \lambda + 1)} \right) + \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})}}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1)} \right\}, \quad (27)
 \end{aligned}$$

3.

$$\|f - Q_{n,m}^*\|_{\infty, [-1,1]^2} \leq \frac{c_{00}}{n^{r-v_1} m^{p-v_2}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \quad (28)$$

where $c_{00} := \dot{C} A_{00}$, with

$$A_{00} := \frac{1}{h_1! h_2!} \left(\sum_{\tau=h_1}^{v_1} \sum_{\mu=h_2}^{v_2} l_{\tau\mu} \frac{2^{(\tau+\mu)-(\alpha_{1\tau}+\alpha_{2\mu})}}{\Gamma(\tau - a_{1\tau} + 1) \Gamma(\mu - \alpha_{2\mu} + 1)} \right) + 1,$$

4.

$$\begin{aligned}
 & \left\| D_{1-}^{(0, \alpha_{2j})} (f) - D_{1-}^{(0, \alpha_{2j})} Q_{n,m}^* \right\|_{\infty, [-1,1]^2} \\
 & \leq \frac{c_{0j}}{n^{r-v_1} m^{p-v_2}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \quad (29)
 \end{aligned}$$

where $c_{0j} = \dot{C} A_{0j}$, $j = 1, \dots, h_2$, with

$$\begin{aligned}
 & A_{0j} \\
 & := \left[\frac{1}{h_1!} \left(\sum_{\tau=h_1}^{v_1} \sum_{\mu=h_2}^{v_2} l_{\tau\mu} \frac{2^{(\tau+\mu)-(\alpha_{1\tau}+\alpha_{2\mu})}}{\Gamma(\tau - a_{1\tau} + 1) \Gamma(\mu - \alpha_{2\mu} + 1)} \right) \right. \\
 & \cdot \left. \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2 - \alpha_{2j} - \lambda + 1)} \right) + \frac{2^{j-\alpha_{2j}}}{\Gamma(j - \alpha_{2j} + 1)} \right], \quad (30)
 \end{aligned}$$

5.

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, 0)}(f) - D_{1-}^{(\alpha_{1i}, 0)} Q_{n,m}^* \right\|_{\infty, [-1, 1]^2} \\ & \leq \frac{c_{i0}}{n^{r-v_1} m^{p-v_2}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \end{aligned} \quad (31)$$

where $c_{i0} = \dot{C} A_{i0}$, $i = i, \dots, h_1$, with

$$\begin{aligned} & A_{i0} \\ & := \left[\frac{1}{h_2!} \left(\sum_{\tau=h_1}^{v_1} \sum_{\mu=h_2}^{v_2} l_{\tau\mu} \frac{2^{(\tau+\mu)-(\alpha_{1\tau}+\alpha_{2\mu})}}{\Gamma(\tau-a_{1\tau}+1) \Gamma(\mu-\alpha_{2\mu}+1)} \right) \right. \\ & \cdot \left. \left(\sum_{k=0}^{h_1-i} \frac{2^{h_1-\alpha_{1i}-k}}{k! \Gamma(h_1-\alpha_{1i}-k+1)} \right) + \frac{2^{i-\alpha_{1i}}}{\Gamma(i-\alpha_{1i}+1)} \right]. \end{aligned} \quad (32)$$

Proof By Corollary 3 there exists

$$Q_{n,m} \equiv Q_{n,m}(f) \in (P_n \otimes C([-1, 1]) + C([-1, 1]) \otimes P_m)$$

such that

$$\|f^{(i,j)} - Q_{n,m}^{(i,j)}\|_{\infty} \leq \frac{\dot{C}}{n^{r-i} m^{p-j}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \quad (33)$$

for all $(0, 0) \leq (i, j) \leq (r, p)$, while $Q_{n,m} \in C^{r,p}([-1, 1])^2$. Here \dot{C} depends only on r, p, s, q , where $n \geq \max\{4(r+1), r+s\}$ and $m \geq \max\{4(p+1), p+q\}$, with $r, p \in \mathbb{Z}_+$, $s, q \in \mathbb{N}$, $f \in C^{r,p}([-1, 1]^2)$. Indeed by [5] we have that $Q_{n,m}^{(r,p)}$ is continuous on $[-1, 1]^2$. We observe the following ($i = 1, \dots, r$; $j = 1, \dots, p$)

$$\begin{aligned} & \left| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f(x_1, x_2) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}(x_1, x_2) \right| \\ & = \frac{1}{\Gamma(i-\alpha_{1i}) \Gamma(j-\alpha_{2j})} \left| \int_{x_1}^1 \int_{x_2}^1 (t_1 - x_1)^{i-\alpha_{1i}-1} (t_2 - x_2)^{j-\alpha_{2j}-1} \right. \\ & \cdot \left. \left(\frac{\partial^{i+j} f(t_1, t_2)}{\partial t_1^i \partial t_2^j} - \frac{\partial^{i+j} Q_{n,m}(t_1, t_2)}{\partial t_1^i \partial t_2^j} \right) dt_1 dt_2 \right| \end{aligned} \quad (34)$$

$$\leq \frac{1}{\Gamma(i-\alpha_{1i}) \Gamma(j-\alpha_{2j})} \int_{x_1}^1 \int_{x_2}^1 (t_1 - x_1)^{i-\alpha_{1i}-1} (t_2 - x_2)^{j-\alpha_{2j}-1} \quad (35)$$

$$\begin{aligned} & \left| \frac{\partial^{i+j} f(t_1, t_2)}{\partial t_1^i \partial t_2^j} - \frac{\partial^{i+j} Q_{n,m}(t_1, t_2)}{\partial t_1^i \partial t_2^j} \right| dt_1 dt_2 \\ & \stackrel{(9)}{\leq} \frac{1}{\Gamma(i - \alpha_{1i}) \Gamma(j - \alpha_{2j})} \\ & \cdot \left(\int_{x_1}^1 \int_{x_2}^1 (t_1 - x_1)^{i - \alpha_{1i} - 1} (t_2 - x_2)^{j - \alpha_{2j} - 1} \right) \\ & \cdot \frac{\dot{C}}{n^{r-i} m^{p-j}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \end{aligned} \tag{36}$$

$$\begin{aligned} & = \frac{1}{\Gamma(i - \alpha_{1i}) \Gamma(j - \alpha_{2j})} \frac{(1 - x_1)^{i - \alpha_{1i}} (1 - x_2)^{j - \alpha_{2j}}}{i - \alpha_{1i} \quad j - \alpha_{2j}} \frac{\dot{C}}{n^{r-i} m^{p-j}} \\ & \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \\ & = \frac{(1 - x_1)^{i - \alpha_{1i}} (1 - x_2)^{j - \alpha_{2j}}}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1)} \frac{\dot{C}}{n^{r-i} m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right). \end{aligned} \tag{37}$$

That is there exists $Q_{n,m}$:

$$\begin{aligned} & \left| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f(x_1, x_2) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}(x_1, x_2) \right| \\ & \leq \frac{(1 - x_1)^{i - \alpha_{1i}} (1 - x_2)^{j - \alpha_{2j}}}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1)} \frac{\dot{C}}{n^{r-i} m^{p-j}} \cdot \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \end{aligned} \tag{38}$$

$i = 1, \dots, r, j = 1, \dots, p, \forall (x_1, x_2) \in [-1, 1]^2$.

We proved there exists $Q_{n,m}$ such that

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} (f) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^* \right\|_{\infty} \\ & \leq \frac{2^{(i+j) - (\alpha_{1i} + \alpha_{2j})} \dot{C}}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1) n^{r-i} m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \end{aligned} \tag{39}$$

$i = 0, 1, \dots, r, j = 0, 1, \dots, p$. Define

$$\rho_{n,m} \equiv \dot{C} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \tag{40}$$

$$\cdot \left[\sum_{i=h_1}^{v_1} \sum_{j=h_2}^{v_2} \left(l_{ij} \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})}}{\Gamma(i-\alpha_{1i}+1)\Gamma(j-\alpha_{2j}+1)} n^{i-r} m^{j-p} \right) \right].$$

I. Suppose, throughout $[-1, 0]^2$, $\alpha_{h_1 h_2}(x, y) \geq \alpha > 0$. Let $Q_{n,m}^*(x, y)$, $(x, y) \in [-1, 1]^2$, as in (39), so that

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} \left(f(x, y) + \rho_{n,m} \frac{x^{h_1} y^{h_2}}{h_1! h_2!} \right) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^*(x, y) \right\|_{\infty} \\ & \leq \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})} \dot{C}}{\Gamma(i-\alpha_{1i}+1)\Gamma(j-\alpha_{2j}+1) n^{r-i} m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \\ & =: T_{ij}, \end{aligned} \quad (41)$$

$i = 0, 1, \dots, r; j = 0, 1, \dots, p$.

If $(h_1 + 1, h_2 + 1) \leq (i, j) \leq (r, p)$, or $0 < i \leq h_1, h_2 + 1 \leq j \leq p$, or $h_1 + 1 \leq i \leq r, 0 < j \leq h_2$ we get from the last

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} (f) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^* \right\|_{\infty} \\ & \leq \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})} \dot{C}}{\Gamma(i-\alpha_{1i}+1)\Gamma(j-\alpha_{2j}+1) n^{r-i} m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \end{aligned} \quad (42)$$

proving (25).

If $(0, 0) < (i, j) \leq (h_1, h_2)$, we get

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f(x, y) + \rho_{n,m} D_{1-}^{\alpha_{1i}} \left(\frac{x^{h_1}}{h_1!} \right) D_{1-}^{\alpha_{2j}} \left(\frac{y^{h_2}}{h_2!} \right) \right. \\ & \left. - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^*(x, y) \right\|_{\infty} \leq T_{ij}. \end{aligned} \quad (43)$$

That is for $i = 1, \dots, h_1; j = 1, \dots, h_2$, we obtain

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f(x, y) + \rho_{n,m} \left((-1)^{h_1} \sum_{k=0}^{h_1-i} \frac{(-1)^k (1-x)^{h_1-\alpha_{1i}-k}}{k! \Gamma(h_1-\alpha_{1i}-k+1)} \right) \right. \\ & \cdot \left((-1)^{h_2} \sum_{\lambda=0}^{h_2-j} \frac{(-1)^\lambda (1-y)^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2-\alpha_{2j}-\lambda+1)} \right) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^*(x, y) \right\|_{\infty} \\ & \leq T_{ij}. \end{aligned} \quad (44)$$

Hence for $(1, 1) \leq (i, j) \leq (h_1, h_2)$, we have

$$\begin{aligned} & \left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^* \right\|_{\infty} \\ & \leq \rho_{n,m} \left(\sum_{k=0}^{h_1-i} \frac{2^{h_1-\alpha_{1i}-k}}{k! \Gamma(h_1 - \alpha_{1i} - k + 1)} \right) \\ & \quad \cdot \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2 - \alpha_{2j} - \lambda + 1)} \right) + T_{ij} \end{aligned} \tag{45}$$

$$= \dot{C} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \tag{46}$$

$$\begin{aligned} & \cdot \left[\sum_{\bar{i}=h_1}^{v_1} \sum_{\bar{j}=h_2}^{v_2} l_{\bar{i}\bar{j}} \frac{2^{(\bar{i}+\bar{j})-(\alpha_{1\bar{i}}+\alpha_{2\bar{j}})}}{\Gamma(\bar{i} - \alpha_{1\bar{i}} + 1) \Gamma(\bar{j} - \alpha_{2\bar{j}} + 1)} \frac{1}{n^{r-\bar{i}}} \frac{1}{m^{p-\bar{j}}} \right] \\ & \left(\sum_{k=0}^{h_1-i} \frac{2^{h_1-\alpha_{1i}-k}}{k! \Gamma(h_1 - \alpha_{1i} - k + 1)} \right) \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2 - \alpha_{2j} - \lambda + 1)} \right) \\ & + \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})} \dot{C} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right)}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1) n^{r-i} m^{p-j}} \\ & \leq \dot{C} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \frac{1}{n^{r-v_1} m^{p-v_2}} A_{ij}, \end{aligned} \tag{47}$$

where

$$\begin{aligned} & A_{ij} \\ & := \left\{ \left[\sum_{\bar{i}=h_1}^{v_1} \sum_{\bar{j}=h_2}^{v_2} l_{\bar{i}\bar{j}} \frac{2^{(\bar{i}+\bar{j})-(\alpha_{1\bar{i}}+\alpha_{2\bar{j}})}}{\Gamma(\bar{i} - \alpha_{1\bar{i}} + 1) \Gamma(\bar{j} - \alpha_{2\bar{j}} + 1)} \right] \right. \\ & \quad \cdot \left(\sum_{k=0}^{h_1-i} \frac{2^{h_1-\alpha_{1i}-k}}{k! \Gamma(h_1 - \alpha_{1i} - k + 1)} \right) \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2 - \alpha_{2j} - \lambda + 1)} \right) \\ & \quad \left. + \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})}}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1)} \right\}. \end{aligned} \tag{48}$$

(Set $c_{ij} := CA_{ij}$)

We proved, for $(1, 1) \leq (i, j) \leq (h_1, h_2)$, that

$$\left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^* \right\|_{\infty} \leq \frac{c_{ij}}{n^{r-v_1} m^{p-v_2}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right). \tag{49}$$

So that (26) is established.

When $i = j = 0$ from (41) we obtain

$$\left\| f(x, y) + \rho_{n,m} \frac{x^{h_1} y^{h_2}}{h_1! h_2!} - Q_{n,m}^*(x, y) \right\|_{\infty} \leq \frac{\dot{C}}{n^r m^p} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right). \quad (50)$$

Hence

$$\|f - Q_{n,m}^*\|_{\infty} \leq \frac{\rho_{n,m}}{h_1! h_2!} + \frac{\dot{C}}{n^r m^p} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \quad (51)$$

$$= \frac{\dot{C}}{h_1! h_2!} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \quad (52)$$

$$\begin{aligned} & \cdot \left[\sum_{\bar{i}=h_1}^{v_1} \sum_{\bar{j}=h_2}^{v_2} l_{\bar{i}\bar{j}} \frac{2^{(\bar{i}+\bar{j})-(\alpha_{1\bar{i}}+\alpha_{2\bar{j}})}}{\Gamma(\bar{i}-\alpha_{1\bar{i}}+1) \Gamma(\bar{j}-\alpha_{2\bar{j}}+1)} \frac{1}{n^{r-\bar{i}}} \frac{1}{m^{p-\bar{j}}} \right] \\ & + \frac{\dot{C}}{n^r m^p} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \\ & \leq \frac{\dot{C} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right)}{n^{r-v_1} m^{p-v_2}} A_{00}, \end{aligned} \quad (53)$$

where

$$A_{00} := \left[\frac{1}{h_1! h_2!} \sum_{\bar{i}=h_1}^{v_1} \sum_{\bar{j}=h_2}^{v_2} \frac{l_{\bar{i}\bar{j}} 2^{(\bar{i}+\bar{j})-(\alpha_{1\bar{i}}+\alpha_{2\bar{j}})}}{\Gamma(\bar{i}-\alpha_{1\bar{i}}+1) \Gamma(\bar{j}-\alpha_{2\bar{j}}+1)} + 1 \right]. \quad (54)$$

(Set $c_{00} = \dot{C} A_{00}$).

Then

$$\|f - Q_{n,m}^*\|_{\infty} \leq \frac{c_{00}}{n^{r-v_1} m^{p-v_2}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right). \quad (55)$$

So that (28) is established.

Next case of $i = 0, j = 1, \dots, h_2$, from (41) we get

$$\left\| D_{1-}^{(0,\alpha_{2j})} f(x, y) + \rho_{n,m} \frac{x^{h_1}}{h_1!} \left((-1)^{h_2} \sum_{\lambda=0}^{h_2-j} \frac{(-1)^\lambda (1-y)^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2-\alpha_{2j}-\lambda+1)} \right) - D_{1-}^{(0,\alpha_{2j})} Q_{n,m}^*(x, y) \right\|_\infty \leq T_{0j}. \tag{56}$$

Then

$$\begin{aligned} & \left\| D_{1-}^{(0,\alpha_{2j})} f - D_{1-}^{(0,\alpha_{2j})} Q_{n,m}^* \right\|_\infty \\ & \leq \frac{\rho_{n,m}}{h_1!} \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2-\alpha_{2j}-\lambda+1)} \right) + T_{0j} \end{aligned} \tag{57}$$

$$= \frac{\dot{C}}{h_1!} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \tag{58}$$

$$\begin{aligned} & \cdot \left[\sum_{\bar{i}=h_1}^{v_1} \sum_{\bar{j}=h_2}^{v_2} l_{\bar{i}\bar{j}} \frac{2^{(\bar{i}+\bar{j})-(\alpha_{1\bar{i}}+\alpha_{2\bar{j}})}}{\Gamma(\bar{i}-\alpha_{1\bar{i}}+1) \Gamma(\bar{j}-\alpha_{2\bar{j}}+1)} \frac{1}{n^{r-\bar{i}}} \frac{1}{m^{p-\bar{j}}} \right] \\ & \cdot \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2-\alpha_{2j}-\lambda+1)} \right) \\ & + \frac{2^{j-\alpha_{2j}} \dot{C}}{\Gamma(j-\alpha_{2j}+1) n^r m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \\ & \leq \frac{\dot{C} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right)}{n^{r-v_1} m^{p-v_2}} A_{0j}, \end{aligned} \tag{59}$$

where

$$\begin{aligned} & A_{0j} \\ & := \left[\frac{1}{h_1!} \left(\sum_{\bar{i}=h_1}^{v_1} \sum_{\bar{j}=h_2}^{v_2} l_{\bar{i}\bar{j}} \frac{2^{(\bar{i}+\bar{j})-(\alpha_{1\bar{i}}+\alpha_{2\bar{j}})}}{\Gamma(\bar{i}-\alpha_{1\bar{i}}+1) \Gamma(\bar{j}-\alpha_{2\bar{j}}+1)} \right) \right. \\ & \cdot \left. \left(\sum_{\lambda=0}^{h_2-j} \frac{2^{h_2-\alpha_{2j}-\lambda}}{\lambda! \Gamma(h_2-\alpha_{2j}-\lambda+1)} \right) + \frac{2^{j-\alpha_{2j}}}{\Gamma(j-\alpha_{2j}+1)} \right]. \end{aligned} \tag{60}$$

(Set $c_{0j} := \dot{C} A_{0j}$)

We proved that (case of $i = 0, j = 1, \dots, h_2$)

$$\left\| D_{1-}^{(0,\alpha_{2j})} f - D_{1-}^{(0,\alpha_{2j})} Q_{n,m}^* \right\|_\infty \leq \frac{c_{0j}}{n^{r-v_1} m^{p-v_2}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right). \tag{61}$$

establishing (29).

Similarly we get for $i = 1, \dots, h_1, j = 0$, that

$$\left\| D_{1-}^{(\alpha_{1i}, 0)} f - D_{1-}^{(\alpha_{1i}, 0)} Q_{n,m}^* \right\|_{\infty} \leq \frac{c_{i0}}{n^{r-v_1} m^{p-v_2}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \quad (62)$$

where $c_{i0} := \dot{C} A_{i0}$, with

$$\begin{aligned} & A_{i0} \\ & := \left[\frac{1}{h_2!} \left(\sum_{\bar{i}=h_1}^{v_1} \sum_{\bar{j}=h_2}^{v_2} l_{\bar{i}\bar{j}} \frac{2^{(\bar{i}+\bar{j})-(\alpha_{1\bar{i}}+\alpha_{2\bar{j}})}}{\Gamma(\bar{i}-\alpha_{1\bar{i}}+1) \Gamma(\bar{j}-\alpha_{2\bar{j}}+1)} \right) \right. \\ & \quad \cdot \left. \left(\sum_{k=0}^{h_1-i} \frac{2^{h_1-\alpha_{1i}-k}}{k! \Gamma(h_1-\alpha_{1i}-k+1)} \right) + \frac{2^{i-\alpha_{1i}}}{\Gamma(i-\alpha_{1i}+1)} \right], \end{aligned} \quad (63)$$

deriving (31). So if $(x, y) \in [-1, 0]^2$, then

$$\alpha_{h_1 h_2}^{-1}(x, y) \bar{L}(Q_{n,m}^*(x, y)) \quad (64)$$

$$\begin{aligned} & = \alpha_{h_1 h_2}^{-1}(x, y) \bar{L}(f(x, y)) + \rho_{n,m} \frac{(1-x)^{h_1-\alpha_{1i}}}{\Gamma(h_1-\alpha_{1i}+1)} \frac{(1-y)^{h_2-\alpha_{2j}}}{\Gamma(h_2-\alpha_{2j}+1)} \\ & + \sum_{i=h_1}^{v_1} \sum_{j=h_2}^{v_2} \alpha_{h_1 h_2}^{-1}(x, y) \alpha_{ij}(x, y) \\ & \cdot \left[D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^*(x, y) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f(x, y) - \rho_{n,m} D_{1-}^{(\alpha_{1i}, \alpha_{2j})} \left(\frac{x^{h_1}}{h_1!} \frac{y^{h_2}}{h_2!} \right) \right] \end{aligned} \quad (65)$$

$$\begin{aligned} & \stackrel{(41)}{\geq} \rho_{n,m} \frac{(1-x)^{h_1-\alpha_{1i}}}{\Gamma(h_1-\alpha_{1i}+1)} \frac{(1-y)^{h_2-\alpha_{2j}}}{\Gamma(h_2-\alpha_{2j}+1)} \\ & - \left[\sum_{i=h_1}^{v_1} \sum_{j=h_2}^{v_2} l_{ij} \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})}}{\Gamma(i-\alpha_{1i}+1) \Gamma(j-\alpha_{2j}+1)} \frac{\dot{C}}{n^{r-i} m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \right] \end{aligned} \quad (66)$$

$$\begin{aligned}
 &= \rho_{n,m} \left[\frac{(1-x)^{h_1-\alpha_{1i}}}{\Gamma(h_1-\alpha_{1i}+1)} \frac{(1-y)^{h_2-\alpha_{2j}}}{\Gamma(h_2-\alpha_{2j}+1)} - 1 \right] \tag{67} \\
 &\geq \rho_{n,m} \left[\frac{1}{\Gamma(h_1-\alpha_{1i}+1) \Gamma(h_2-\alpha_{2j}+1)} - 1 \right]
 \end{aligned}$$

$$= \rho_{n,m} \left[\frac{1 - \Gamma(h_1 - \alpha_{1i} + 1) \Gamma(h_2 - \alpha_{2j} + 1)}{\Gamma(h_1 - \alpha_{1i} + 1) \Gamma(h_2 - \alpha_{2j} + 1)} \right] \geq 0. \tag{68}$$

Explanation: we have that $\Gamma(1) = 1$, $\Gamma(2) = 1$, and Γ is convex on $(0, \infty)$ and positive there, here $0 \leq h_1 - \alpha_{1h_1}, h_2 - \alpha_{2h_2} < 1$ and $1 \leq h_1 - \alpha_{1h_1} + 1, h_2 - \alpha_{2h_2} + 1 < 2$. Thus $0 < \Gamma(h_1 - \alpha_{1h_1} + 1), \Gamma(h_2 - \alpha_{2h_2} + 1) \leq 1$, and

$$0 \leq \Gamma(h_1 - \alpha_{1h_1} + 1) \Gamma(h_2 - \alpha_{2h_2} + 1) \leq 1. \tag{69}$$

And

$$1 - \Gamma(h_1 - \alpha_{1h_1} + 1) \Gamma(h_2 - \alpha_{2h_2} + 1) \geq 0. \tag{70}$$

Therefore it holds

$$\bar{L}(Q_{n,m}^*(x, y)) \geq 0, \forall (x, y) \in [-1, 0]^2. \tag{71}$$

II. Suppose, throughout $[-1, 0]^2$, $\alpha_{h_1 h_2}(x, y) \leq \beta < 0$. Let $Q_{n,m}^{**}(x, y), (x, y) \in [-1, 1]^2$, as in (39), so that

$$\begin{aligned}
 &\left\| D_{1-}^{(\alpha_{1i}, \alpha_{2j})} \left(f(x, y) - \rho_{n,m} \frac{x^{h_1}}{h_1!} \frac{y^{h_2}}{h_2!} \right) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^{**}(x, y) \right\|_{\infty} \\
 &\leq \frac{2^{(i+j)-(\alpha_{1i}+\alpha_{2j})} \dot{C}}{\Gamma(i-\alpha_{1i}+1) \Gamma(j-\alpha_{2j}+1) n^{r-i} m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right), \tag{72}
 \end{aligned}$$

$i = 0, 1, \dots, r, j = 0, 1, \dots, p$.

As earlier we produce the same convergence inequalities (25), (26), (28), (29), and (31). So for $(x, y) \in [-1, 0]^2$ we get

$$\begin{aligned} & \alpha_{h_1 h_2}^{-1}(x, y) \bar{L}(Q_{n,m}^{**}(x, y)) \\ &= \alpha_{h_1 h_2}^{-1}(x, y) \bar{L}(f(x, y)) - \rho_{n,m} \frac{(1-x)^{h_1 - \alpha_{1i}}}{\Gamma(h_1 - \alpha_{1i} + 1)} \frac{(1-y)^{h_2 - \alpha_{2j}}}{\Gamma(h_2 - \alpha_{2j} + 1)} \end{aligned} \quad (73)$$

$$\begin{aligned} & + \sum_{i=h_1}^{v_1} \sum_{j=h_2}^{v_2} \alpha_{h_1 h_2}^{-1}(x, y) \alpha_{ij}(x, y) \\ & \cdot \left[D_{1-}^{(\alpha_{1i}, \alpha_{2j})} Q_{n,m}^{**}(x, y) - D_{1-}^{(\alpha_{1i}, \alpha_{2j})} f(x, y) + \rho_{n,m} D_{1-}^{(\alpha_{1i}, \alpha_{2j})} \left(\frac{x^{h_1}}{h_1!} \frac{y^{h_2}}{h_2!} \right) \right] \\ & \stackrel{(71)}{\leq} -\rho_{n,m} \frac{(1-x)^{h_1 - \alpha_{1i}}}{\Gamma(h_1 - \alpha_{1i} + 1)} \frac{(1-y)^{h_2 - \alpha_{2j}}}{\Gamma(h_2 - \alpha_{2j} + 1)} \quad (74) \\ & + \left[\sum_{i=h_1}^{v_1} \sum_{j=h_2}^{v_2} l_{ij} \frac{2^{(i+j) - (\alpha_{1i} + \alpha_{2j})}}{\Gamma(i - \alpha_{1i} + 1) \Gamma(j - \alpha_{2j} + 1)} \frac{\dot{C}}{n^{r-i} m^{p-j}} \omega_{s,q} \left(f^{(r,p)}; \frac{1}{n}, \frac{1}{m} \right) \right] \\ &= \rho_{n,m} \left[1 - \frac{(1-x)^{h_1 - \alpha_{1i}}}{\Gamma(h_1 - \alpha_{1i} + 1)} \frac{(1-y)^{h_2 - \alpha_{2j}}}{\Gamma(h_2 - \alpha_{2j} + 1)} \right] \\ &= \rho_{n,m} \left[\frac{\Gamma(h_1 - \alpha_{1i} + 1) \Gamma(h_2 - \alpha_{2j} + 1) - (1-x)^{h_1 - \alpha_{1i}} (1-y)^{h_2 - \alpha_{2j}}}{\Gamma(h_1 - \alpha_{1i} + 1) \Gamma(h_2 - \alpha_{2j} + 1)} \right] \\ &\leq \rho_{n,m} \left[\frac{1 - (1-x)^{h_1 - \alpha_{1i}} (1-y)^{h_2 - \alpha_{2j}}}{\Gamma(h_1 - \alpha_{1i} + 1) \Gamma(h_2 - \alpha_{2j} + 1)} \right] \leq 0. \quad (75) \end{aligned}$$

Explanation: for $x, y \in [-1, 0]$ we get that $1 - x, 1 - y \geq 1$, and $0 \leq h_1 - \alpha_{1h_1}, h_2 - \alpha_{2h_2} < 1$. Hence $(1-x)^{h_1 - \alpha_{1h_1}}, (1-y)^{h_2 - \alpha_{2h_2}} \geq 1$, and then

$$(1-x)^{h_1 - \alpha_{1h_1}} (1-y)^{h_2 - \alpha_{2h_2}} \geq 1,$$

so that

$$1 - (1-x)^{h_1 - \alpha_{1h_1}} (1-y)^{h_2 - \alpha_{2h_2}} \leq 0. \quad (76)$$

Hence again

$$\bar{L}(Q_{n,m}^{**}(x, y)) \geq 0, \text{ for } (x, y) \in [-1, 0]^2. \quad (77)$$

References

1. Anastassiou, G.A.: Monotone approximation by pseudopolynomials. In: Approximation Theory. Academic Press, New York (1991)
2. Anastassiou, G.A.: Bivariate monotone approximation. Proc. Amer. Math. Soc. **112**(4), 959–963 (1991)

3. Anastassiou, G.A., Shisha, O.: Monotone approximation with linear differential operators. *J. Approx. Theory* **44**, 391–393 (1985)
4. Gonska, H.H., Simultaneously approximation by algebraic blending functions. In: Alfred Haar Memorial Conference, Budapest, *Coloquia Mathematica Soc. Janos Bolyai*, vol. 49, pp. 363–382. North-Holand, Amsterdam (1985)
5. Gonska, H.H.: Personal communication with author, 2–24–2014
6. S. Iqbal, K. Krulic, J. Pecaric, On an inequality of H.G. Hardy, *J. Inequal. Appl.* Article ID 264347, 23pp. (2010)
7. A. Marchaud, Differences et deerivees d'une fonction de deux variables. *C.R. Acad. Sci.* **178**, 1467–1470 (1924)
8. Marchaud, A.: Sur les derivees et sur les differences des fonctions de variables reelles. *J. Math. Pures Appl.* **6**, 337–425 (1927)
9. Schumaker, L.L.: *Spline Functions: Basic Theory*. Wiley, New York (1981)
10. Shisha, O.: Monotone approximation. *Pacific J. Math.* **15**, 667–671 (1965)



<http://www.springer.com/978-3-319-30320-8>

Intelligent Mathematics II: Applied Mathematics and
Approximation Theory

Anastassiou, G.A.; Duman, O. (Eds.)

2016, XV, 502 p. 54 illus., 45 illus. in color., Softcover

ISBN: 978-3-319-30320-8