

The overall age-adjusted incidence of mycosis fungoides (MF) worldwide is around 6–7 cases per 1 million [1]. MF is thought to be a disease of the elderly, as 75 % of cases are seen in the age group of 50–60 years [2]. However, MF can be seen in patients under the age of 35 years old with similar findings and clinical course [3]. MF has in fact been identified in children and teenagers [4]; 0.5–5 % of all cases are diagnosed before age 20 years [5, 6]. In addition, MF is observed more commonly in males than females, with a male-to-female ratio of 2:1 [5, 7]—a finding that is more pronounced in children. In a study looking at juvenile-onset MF, 24 out of 34 patients were male (74 %) [1].

In the United States, the incidence of MF, up until the year 2000, was reported to have increased over time [8], a phenomenon that may be reflective of improvements in diagnostics or reporting. Between 1973 and 2002, an increase in the incidence of MF was found to correlate with a higher density of medical specialists. However according to the Surveillance, Epidemiology, and End Results (SEER) registry data, the incidence of MF appears to have stabilized since then [9].

---

**Contributors to This Chapter**

- Pooya Khan Mohammad Beigi, MD, University of British Columbia, BC, Canada
- Elizabeth Alice Wang, BS, University of California Davis, Sacramento, CA, USA
- Seyed Sajad Niyiyati, BS, University of British Columbia, Vancouver, BC, Canada
- Emanuel Maverakis, MD, University of California Davis, Sacramento, CA, USA

## References

1. Hall JC, Hall BJ. Cutaneous lymphoma: diagnosis and treatment. Shelton: PMPH; 2012.
2. Fatemi Nf, Najafian J, Salehi M, Azimi Z, Nilforoushzadeh Ma, Rajabi P. Mycosis fungoides: epidemiology in Isfahan, Iran; 2014.
3. Crowley JJ, Nikko A, Varghese A, Hoppe RT, Kim YH. Mycosis fungoides in young patients: clinical characteristics and outcome. *J Am Acad Dermatol.* 1998;38(5):696–701.
4. Kim ST, Sim HJ, Jeon YS, Lee JW, Roh HJ, Choi SY, et al. Clinicopathological features and t-cell receptor gene rearrangement findings of mycosis fungoides in patients younger than age 20 years. *J Dermatol.* 2009;36(7):392–402.
5. Weinstock MA, Horm JW. Mycosis fungoides in the United States: increasing incidence and descriptive epidemiology. *JAMA.* 1988;260(1):42–6.
6. Agnarsson BA, Kadin ME, editors. Peripheral t-cell lymphomas in children. *Seminars in diagnostic pathology*; 1995.
7. Willemze R, Jaffe ES, Burg G, Cerroni L, Berti E, Swerdlow SH, et al. WHO-EORTC classification for cutaneous lymphomas. *Blood.* 2005;105(10):3768–85.
8. Morales Suarez-Varela MM, Llopis Gonzalez A, Marquina Vila A, Bell J. Mycosis fungoides: review of epidemiological observations. *Dermatology (Basel, Switzerland).* 2000;201(1): 21–8.
9. Korgavkar K, Xiong M, Weinstock M. Changing incidence trends of cutaneous T-cell lymphoma. *JAMA Dermatol.* 2013;149(11):1295–9.



<http://www.springer.com/978-3-319-47906-4>

Clinician's Guide to Mycosis Fungoides

Khan Mohammad Beigi, P.

2017, XV, 139 p. 91 illus., 86 illus. in color., Hardcover

ISBN: 978-3-319-47906-4