

Contents

Editorial Preface		xi
Wave Maps and Ill-posedness of their Cauchy Problem		
<i>Piero D’Ancona and Vladimir Georgiev</i>		1
1 Preface		1
2 Variational motivation of the wave maps equations		4
2.1 Introduction		4
2.2 Harmonic maps and special harmonic maps on the sphere		6
2.3 Equivariant wave maps and construction of special solutions		12
3 Local existence result for equivariant wave maps		17
3.1 Introduction		17
3.2 Localization in time		20
3.3 Estimates for the homogeneous problem		24
3.4 Estimates for the non-homogeneous problem		26
3.5 Bilinear estimates for the homogeneous problem in $H^{s,\delta}$		31
3.6 Bilinear estimates in $H^{s,\delta}$ for the inhomogeneous problem		36
4 Concentration of the local energy		40
4.1 Introduction		40
4.2 Construction of the solutions		42
4.3 Higher regularity of the solution		46
4.4 Appendix		52
5 Non-uniqueness result in the subcritical case		55
5.1 Introduction		55
5.2 Equivariant and self-similar solutions		57
5.3 Low regularity self-similar solutions		60
5.4 Appendix A: The self-similar ODE		66
5.5 Appendix B: Some technical lemmas		72
6 Ill-posedness in the critical case (Fourier analysis approach)		77
6.1 Introduction		77
6.2 Well-posedness of the Cauchy problem for semilinear wave equations		79

6.3	The wave map system in stereographic projection	80
6.4	Conclusion of the proof of Theorem 6.1	83
6.5	Proof of Theorem 6.2	86
7	Ill-posedness in the critical case (fundamental solution approach)	90
7.1	Introduction	90
7.2	Proof of Theorem 7.1	92
7.3	Appendix	102
	References	109

On the Global Behavior of Classical Solutions to Coupled Systems of Semilinear Wave Equations

	<i>Hideo Kubo and Masahito Ohta</i>	113
1	Introduction	113
2	Single wave equation	117
2.1	Blow-up	119
2.2	Small data global existence	125
2.3	Almost global existence	133
2.4	Self-similar solution	137
2.5	Asymptotic behavior	140
3	Semilinear system, I	156
3.1	Blow-up	157
3.2	Small data global existence	163
3.3	Self-similar solution	166
3.4	Asymptotic behavior	167
4	Semilinear system, II	168
4.1	Small data global existence	170
4.2	Self-similar solution	173
4.3	Generalization	176
5	Semilinear system, III	178
5.1	Blow-up	179
5.2	Small data global existence	183
6	Nonlinear system	187
6.1	Blow-up	192
6.2	Null condition	196
	Appendix	202
	References	205

Decay and Global Existence for Nonlinear Wave Equations with Localized Dissipations in General Exterior Domains

	<i>Mitsuhiro Nakao</i>	213
1	Introduction	213
2	Preliminaries	216

3	Local energy decay	218
3.1	Problem and result	218
3.2	Proof of Theorem 3.1.	219
3.3	Proof of Corollary 3.1.	223
4	Total Energy decay for the wave equation with a localized dissipation	226
4.1	Problem and result	226
4.2	Proof of Theorem 4.1	227
4.3	Proof of Theorem 4.2	232
5	Linear equations with variable coefficients; Unique continuation property and a basic inequality	233
5.1	Problem and result	233
5.2	Proof of Proposition 5.1	235
5.3	Proof of Theorems 5.1 and 5.2	237
5.4	Proof of Proposition 5.2	239
6	L^p estimates for the wave equation in exterior domains	241
6.1	Problem and result	241
6.2	Proof of Theorem 6.2	243
7	Semilinear wave equations	249
7.1	Problem and result	249
7.2	Proof of Theorems 7.1 and 7.2	251
7.3	Proof of Theorem 7.3	253
8	Quasilinear wave equations	259
8.1	Problem and result	259
8.2	Energy decay for the quasilinear equation	261
8.3	Estimation of higher-order derivatives of solutions	265
8.4	Proof of Theorems 8.2 and 8.3.	272
9	The wave equation with a half-linear dissipation	280
9.1	Problem and result	280
9.2	A basic inequality	283
9.3	Proof of Theorem 9.1	286
9.4	Proof of Theorems 9.2 and 9.3	289
10	Some open problems	293
	References	295

Global Existence in the Cauchy Problem for Nonlinear Wave Equations with Variable Speed of Propagation

	<i>Karen Yagdjian</i>	301
1	Introduction	301
2	Counterexamples to the global existence	303
3	Blow-up for the problem with large potential energy of nonlinearity	320
4	Parametric resonance and wave map type equations	324

5	Proof of Theorem 4.1: Parametric resonance	327
5.1	Some properties of the Hill's equation	327
5.2	Borg's Theorem	334
5.3	Construction of an exponentially increasing solution to Hill's equation	338
5.4	Construction of blow-up solutions	342
6	Coefficient stabilizing to a periodic one. Parametric resonance dominates	344
7	Proof of Theorem 6.1: Perturbation theory	345
8	Nonexistence for equations with permanently restricted domain of influence	354
9	Global existence for a model equation with a polynomially growing coefficient	358
10	An example with an exponentially growing coefficient	362
11	Fast oscillating coefficients: no resonance ?!	370
12	Linear wave equations with oscillating coefficients	373
	References	382

On the Nonlinear Cauchy Problem

	<i>Massimo Cicognani and Luisa Zanghirati</i>	387
1	Introduction	387
2	Well-posedness in C^∞	392
2.1	Function and symbol spaces	392
2.2	Levi conditions	394
2.3	Factorization	397
2.4	The linear problem	398
2.5	Commutators	403
2.6	The equivalent quasilinear system	407
2.7	Local C^∞ solutions	409
2.8	Analytic regularity	410
3	Well-posedness in Gevrey classes	416
3.1	The linear problem	416
3.2	Gevrey-Levi conditions	419
3.3	Factorization under Gevrey-Levi conditions	422
3.4	Linear systems	423
3.5	The equivalent quasilinear system in Gevrey spaces	427
3.6	Local Gevrey solutions and propagation of the analytic regularity	428
4	Strictly hyperbolic equations with non-Lipschitz coefficients and C^∞ solutions	430
4.1	Log-Lipschitz coefficients or unbounded derivatives	430
4.2	The linear problem with non-regular coefficients	432
4.3	The map $u \mapsto v$	436

5	Hölder coefficients and Gevrey Solutions	441
5.1	Gevrey well-posedness	441
5.2	From the factorization to the quasilinear system	442
	References	445

Sharp Energy Estimates for a Class of Weakly Hyperbolic Operators

Michael Dreher and Ingo Witt **449**

1	Introduction	449
1.1	Well-posedness of the Cauchy problem	450
1.2	Degenerate differential operators	451
1.3	Notation	454
2	Formulation of the results	455
2.1	Motivation and plan of the paper	455
2.2	Main results	457
3	A model case	462
3.1	Taniguchi–Tozaki’s example	463
3.2	Conversion into a 2×2 system	464
3.3	Estimation of the fundamental matrix	464
3.4	Function spaces: An approach via edge Sobolev spaces	465
3.5	Establishing energy estimates	470
3.6	Summary of Section 3	470
4	Symbol classes and function spaces	471
4.1	The symbol classes $S^{m,\eta;\lambda}$	471
4.2	The symbol classes $\tilde{S}^{m,\eta;\lambda}$	472
4.3	The symbol classes $S_+^{m,\eta;\lambda}$ for $\eta \in C_b^\infty(\mathbb{R}^n; \mathbb{R})$	474
4.4	Function spaces: An approach via weight functions	475
4.5	Summary of Section 4	478
5	The Cauchy problem	478
5.1	Improvement of Gårding’s inequality	480
5.2	Symmetric-hyperbolic systems	481
5.3	Symmetrizable-hyperbolic systems	483
5.4	Higher-order scalar equations	487
5.5	Local uniqueness	489
5.6	Sharpness of energy estimates	491
A	Supplements	498
B	Open problems	506
	References	507



<http://www.springer.com/978-3-7643-7283-5>

New Trends in the Theory of Hyperbolic Equations

Reissig, M.; Schulze, B.-W. (Eds.)

2005, XIII, 514 p., Hardcover

ISBN: 978-3-7643-7283-5

A product of Birkhäuser Basel